

Number of Years Lost Before The Age of 75 by Cause of Death and Sex, Norrbotten

Indicator	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
All causes, years lost before 75, men (5-year mean value) (1-årsvärde)	95.49	91.31	90.59	89.01	86.32	83.34	82.49	80.35	77.96	76.77	74.24	72.24	69	67.91	65.83
Infectious diseases etc, years lost before 75, men (5-year mean value) (1-årsvärde)	0.43	0.48	0.64	0.6	0.62	0.61	0.72	0.67	0.84	0.79	0.78	0.81	0.72	0.54	0.5
Tumours, years lost before 75, men (5-year mean value) (1-årsvärde)	15.6	14.7	14.48	14.33	14.04	14.49	14.8	14.22	13.93	13.87	12.97	12.86	12.84	12.4	12.53
Cancer of the stomach, years lost before 75, men (5-year mean value) (1-årsvärde)	1.4	1.22	1.41	1.42	1.37	1.49	1.41	1.35	1.12	1.27	1.02	0.94	0.94	0.97	0.86
Cancer of the large intestine, years lost before 75, men (5-year mean value) (1-årsvärde)	0.82	0.77	0.76	0.88	0.92	0.94	0.87	0.84	0.97	0.79	0.65	0.64	0.81	0.63	0.74
Cancer of the liver, biliary passages, years lost before 75, men (5-year mean value) (1-årsvärde)	0.44	0.44	0.3	0.36	0.4	0.43	0.43	0.45	0.37	0.28	0.32	0.27	0.32	0.32	0.55
Lung cancer, years lost before 75, men (5-year mean value) (1-årsvärde)	2.76	2.64	2.69	2.7	2.8	2.71	2.76	2.62	2.55	2.38	2.38	2.34	2.38	2.23	2.16
Malignant melanoma, years lost before 75, men (5-year mean value) (1-årsvärde)	0.27	0.23	0.28	0.33	0.28	0.25	0.28	0.27	0.28	0.26	0.32	0.43	0.34	0.3	0.35
Prostate cancer, years lost before 75, men (5-year mean value) (1-årsvärde)	1.07	1.07	1.2	1.05	1.07	1.15	1.01	0.86	0.95	0.9	0.89	1.05	0.99	0.97	1.06
Cancer of the bladder, not kidneys, years lost before 75, men (5-year mean value) (1-årsvärde)	0.47	0.46	0.39	0.42	0.32	0.2	0.15	0.18	0.18	0.22	0.23	0.22	0.23	0.19	0.2
Leukaemia, years lost before 75, men (5-year mean value) (1-årsvärde)	1.02	0.98	0.72	0.69	0.66	0.53	0.71	0.89	0.95	1.11	1.06	0.95	0.75	0.61	0.34
Endocrine diseases etc, years lost before 75, men (5-year mean value) (1-årsvärde)	1.41	1.18	1.24	1.64	1.93	1.83	2.39	2.42	2.45	2.53	2.88	2.4	2.47	2.12	1.86
Diabetes, years lost before 75, men (5-year mean value) (1-årsvärde)	4.67	4.74	4.91	5.11	5.15	5.28	5.52	5.72	5.3	5.1	4.87	4.49	4.46	4.38	4.47
Mental disorders, years lost before 75, men (5-year mean value) (1-årsvärde)	2.37	2.68	3.34	3.34	3.18	3.07	3.1	2.77	2.66	2.76	2.79	2.52	2.42	2.45	2.58
Diseases of the circulatory organs, years lost before 75, men (5-year mean value) (1-årsvärde)	39.63	38.1	37.84	36.19	34.86	32.62	30.48	29.68	28.01	27.02	26.22	25.78	23.75	23.65	22.16
Ischaemic heart disease, years lost before 75, men (5-year mean value) (1-årsvärde)	30.33	28.62	28.46	27.56	26.09	24.07	22.14	20.64	19.26	18.2	16.98	16.67	15.33	15.07	14
Acute myocardial infarction, years lost before 75, men (5-year mean value) (1-årsvärde)	22.07	20.22	19.86	19.37	18.01	16.18	14.77	13.25	12.37	12.03	11.81	11.72	10.95	10.55	9.43
Stroke, years lost before 75, men (5-year mean value) (1-årsvärde)	4.76	4.7	4.63	3.97	4.02	3.95	3.61	3.83	3.68	3.92	4.18	4.22	3.71	3.94	3.68
Diseases of the respiratory organs, years lost before 75, men (5-year mean value) (1-årsvärde)	2.19	2.19	2.22	2.71	2.69	2.38	2.29	2.3	2.09	1.85	1.87	1.74	1.55	1.32	1.56
Bronchitis, emphysema, asthma, years lost before 75, men (5-year mean value) (1-årsvärde)	1.25	1.01	0.95	1.25	1.26	0.99	1.04	1.24	1.1	0.92	1.01	0.9	0.7	0.57	0.72
Asthma, years lost before 75, men (5-year mean value) (1-årsvärde)	0.52	0.37	0.35	0.54	0.6	0.52	0.54	0.6	0.4	0.29	0.29	0.22	0.12	0.09	0.18
Diseases of the digestive organs, years lost before 75, men (5-year mean value) (1-årsvärde)	2.85	3.17	2.93	2.54	2.37	2.21	1.74	1.73	1.85	1.82	1.83	1.67	1.64	1.49	
Cirrhosis of the liver + other alc. rel. liver diseases, years lost before 75, men (5-year mean value) (1-årsvärde)	1.86	1.92	1.76	1.4	1.23	1.12	0.8	0.75	0.75	0.65	0.67	0.76	0.74	0.66	0.69
Injuries and poisoning, years lost before 75, men (5-year mean value) (1-årsvärde)	27.26	25.56	24.53	24.14	23.28	22.7	23.52	22.66	22.25	21.77	20.7	19.96	19.59	19.69	19.19
Accidents, years lost before 75, men (5-year mean value) (1-årsvärde)	15.02	14.38	14.18	13.41	12.83	12.72	13.05	12.15	12.61	12.02	10.81	10.48	9.85	9.76	9.35
Motor-vehicle accidents, years lost before 75, men (5-year mean value) (1-årsvärde)	6.29	6.3	6.47	6.62	6.54	6.22	6.36	5.86	6.22	5.73	4.9	4.7	4.5	4.51	4.66
Falls, years lost before 75, men (5-year mean value) (1-årsvärde)	1.42	1.19	1.15	1.1	0.86	0.91	0.88	0.78	0.8	0.97	0.81	0.97	0.98	0.95	0.91
Suicide or undetermined intent, years lost before 75, men (5-year mean value) (1-årsvärde)	11.33	10.44	9.86	10.19	9.9	9.59	10.05	9.81	9.02	9.27	9.34	8.92	9.3	9.42	9.31
Murder, manslaughter and other abuse, years lost before 75, men (5-year mean value) (1-årsvärde)	0.88	0.71	0.46	0.55	0.55	0.35	0.38	0.66	0.57	0.44	0.55	0.56	0.44	0.5	0.53
Alcohol-related diseases, years lost before 75, men (5-year mean value) (1-årsvärde)	9.77	9.51	9.31	9.37	9.01	8.47	8.12	7.67	6.89	6.3	6.34	6.13	6.05	6.04	6.26
All causes, years lost before 75, women (5-year mean value) (1-årsvärde)	44.73	42.27	41.22	40.48	40.25	38.99	39.04	39.38	39.02	38.46	36.9	37.21	36.16	34.55	33.75
Infectious diseases etc, years lost before 75, women (5-year mean value) (1-årsvärde)	0.52	0.7	0.58	0.57	0.55	0.69	0.5	0.44	0.5	0.54	0.36	0.35	0.34	0.33	0.35
Tumours, years lost before 75, women (5-year mean value) (1-årsvärde)	15.5	15	14.97	15.25	15.4	14.91	15.14	15.79	15.73	16.14	15.65	15.77	15.09	14.02	13.85
Cancer of the stomach, years lost before 75, women (5-year mean value) (1-årsvärde)	1.14	1.06	0.93	0.92	0.96	0.86	0.87	0.85	0.93	0.83	0.77	0.79	0.72	0.53	0.51
Cancer of the large intestine, years lost before 75, women (5-year mean value) (1-årsvärde)	0.94	1.01	0.88	0.69	0.58	0.54	0.51	0.65	0.85	0.83	0.81	0.96	0.98	0.81	0.91
Lung cancer, years lost before 75, women (5-year mean value) (1-årsvärde)	1.03	1.07	1.44	1.46	1.46	1.55	1.64	1.5	1.49	1.52	1.67	1.67	1.64	1.57	1.65
Malignant melanoma, years lost before 75, women (5-year mean value) (1-årsvärde)	0.31	0.25	0.18	0.18	0.18	0.09	0.09	0.08	0.13	0.12	0.13	0.19	0.23	0.16	0.17
Breast cancer, years lost before 75, women (5-year mean value) (1-årsvärde)	3.07	2.86	2.97	3.07	3.12	2.95	3.11	3.21	3.25	3.52	3.4	3.31	3.22	2.97	2.53
Cervical cancer, years lost before 75, women (5-year mean value) (1-årsvärde)	0.72	0.6	0.63	0.74	0.64	0.44	0.38	0.36	0.17	0.26	0.3	0.31	0.31	0.36	0.26
Ovarian cancer, years lost before 75, women (5-year mean value) (1-årsvärde)	1.23	1.28	1.3	1.55	1.5	1.75	1.77	1.86	1.84	2	1.8	1.85	1.66	1.6	1.35
Cancer of the bladder, not kidneys, years lost before 75, women (5-year mean value) (1-årsvärde)	0.12	0.1	0.11	0.14	0.11	0.1	0.11	0.16	0.19	0.19	0.18	0.18	0.1	0.05	0.05
Leukaemia, years lost before 75, women (5-year mean value) (1-årsvärde)	0.77	0.7	0.77	0.98	1.03	0.66	0.7	0.77	0.75	0.79	0.8	0.73	0.59	0.32	0.22
Endocrine diseases etc, years lost before 75, women (5-year mean value) (1-årsvärde)	1.37	1.41	1.32	1.46	1.34	1.23	1.09	1.06	1	0.89	0.88	1.03	1.08	1.23	1.05
Diabetes, years lost before 75, women (5-year mean value) (1-årsvärde)	3.45	3.35	3.21	3.02	3.02	2.8	2.38	2.21	2.22	1.94	1.84	1.95	2.12	2.04	2.09
Mental disorders, years lost before 75, women (5-year mean value) (1-årsvärde)	0.61	0.77	0.86	0.59	0.7	0.64	0.56	0.62	0.7	0.72	0.75	0.68	0.69	0.7	0.54
Diseases of the circulatory organs, years lost before 75, women (5-year mean value) (1-årsvärde)	13.67	13.12	12.61	11.62	10.62	9.8	9.63	9.33	9.29	9.25	8.99	8.89	8.42	7.95	7.81
Ischaemic heart disease, years lost before 75, women (5-year mean value) (1-årsvärde)	6.99	6.88	6.57	6.3	5.5	5.25	4.93	4.94	4.71	4.97	4.67	4.45	3.94	3.67	3.12
Acute myocardial infarction, years lost before 75, women (5-year mean value) (1-årsvärde)	5.49	5.46	5.04	4.92	4.06	3.74	3.55	3.55	3.26	3.47	3.28	3.23	2.9	2.66	2.16
Stroke, years lost before 75, women (5-year mean value) (1-årsvärde)	4.52	4.15	3.95	3.33	3.02	2.53	2.58	2.39	2.49	2.38	2.52	2.32	2.22	2.21	2.29
Diseases of the respiratory organs, years lost before 75, women (5-year mean value) (1-årsvärde)	1.52	1.53	1.5	1.62	1.69	1.75	1.66	1.68	1.49	1.23	1.21	1.2	1.11	1.14	1.18
Bronchitis, emphysema, asthma, years lost before 75, women (5-year mean value) (1-årsvärde)	0.9	0.97	0.94	1.06	1.21	1.3	1.2	1.23	1.08	0.87	0.81	0.73	0.62	0.68	0.74
Asthma, years lost before 75, women (5-year mean value) (1-årsvärde)	0.64	0.57	0.47	0.5	0.64	0.82	0.84	0.92	0.81	0.51	0.45	0.3	0.23	0.26	0.22
Diseases of the digestive organs, years lost before 75, women (5-year mean value) (1-årsvärde)	1.56	1.51	1.29	1.05	0.94	0.85	0.84	0.97	1.02	0.92	0.78	0.79	0.76	0.64	0.79
Cirrhosis of the liver + other alc. rel. liver diseases, years lost before 75, women (5-year mean value) (1-årsvärde)	0.71	0.57	0.54	0.45	0.51	0.38	0.31	0.37	0.32	0.19	0.18	0.3	0.27	0.26	0.36
Injuries and poisoning, years lost before 75, women (5-year mean value) (1-årsvärde)	7.52	6.21	6.13	6.11	6.72	6.2	6.41	6.17	6.18	5.45	5.33	5.54	5.85	5.92	5.62
Accidents, years lost before 75, women (5-year mean value) (1-årsvärde)	3.86	2.79	2.88	3.03	3.43	3.28	3.4	3.21	3.19	2.49	2.31	2.67	2.78	2.92	2.99
Motor-vehicle accidents, years lost before 75, women (5-year mean value) (1-årsvärde)	2.29	1.62	1.72	2.03	2.23	2.03	2.23	2.36	2.09	1.74	1.59	1.6	1.46	1.71	1.75
Falls, years lost before 75, women (5-year mean value) (1-årsvärde)	0.18	0.15	0.12	0.08	0.13	0.09	0.07	0.06	0.08	0.03	0.03	0.03	0.04	0.07	0.11
Suicide or undetermined intent, years lost before 75, women (5-year mean value) (1-årsvärde)	3.49	3.19	2.97	2.75	2.86	2.54	2.72	2.77	2.81	2.92	2.81	2.61	2.77	2.73	2.26
Murder, manslaughter and other abuse, years lost before 75, women (5-year mean value) (1-årsvärde)	0.17	0.23	0.27	0.32	0.42	0.38	0.28	0.19	0.18	0.04	0.22	0.26	0.31	0.26	0.29
Alcohol-related diseases, years lost before 75, women (5-year mean value) (1-årsvärde)	1.27	1.23	1.11	0.9	1.16	1.05	0.97	1.16	1.18	1.01	1.09	1.11	1.11	1.22	1.29