

Alcohol Use – Binge Drinking

	2002				2003			
	n	weighted %	N	95% CI	n	weighted %	N	95% CI
Gender								
Male	280	25%	1,159	21.7–29.0	321	25%	1,208	21.5–28.1
Female	178	11%	1,489	8.9–13.3	166	12%	1,420	9.3–14.5
Total	458	18%	2,648	16.1–20.6	487	18%	2,628	16.3–20.6
Race								
Native (any mention)	120	20%	572	15.2–25.8	124	24%	562	18.6–29.8
Non-Native	332	18%	2,050	15.5–20.4	359	17%	2,052	15.0–19.6
Age								
18–24	64	25%	243	18.5–32.3	51	23%	221	16.4–31.5
25–34	118	26%	491	19.9–32.4	131	26%	486	20.4–31.5
35–44	123	19%	650	15.2–24.2	134	19%	612	15.2–23.3
45–54	94	15%	623	11.3–19.0	108	18%	644	14.0–22.9
55–64	35	11%	345	7.2–17.3	44	12%	366	8.3–17.6
65 or older	22	6%	282	3.3–11.5	18	6%	277	3.0–11.0
Education								
Some H.S.	43	28%	224	18.3–39.5	45	21%	216	14.3–30.3
H.S. grad or GED	171	21%	862	16.7–24.9	172	21%	814	17.8–25.7
Some College or Tech School	130	16%	776	13.1–20.5	160	20%	812	16.3–24.8
College Grad	114	15%	783	12.0–19.3	108	12%	780	9.5–15.5
Income								
Less than \$15K	38	18%	240	11.7–25.8	45	22%	214	14.4–32.3
\$15,000–24,999	75	21%	389	14.8–28.0	79	26%	360	19.2–33.5
\$25,000–34,999	57	16%	343	10.9–21.6	73	19%	352	13.7–24.6
\$35,000–49,999	69	15%	435	10.6–21.3	76	17%	472	12.8–22.7
\$50,000–74,999	100	25%	455	19.4–31.4	90	17%	468	12.8–21.8
\$75K+	89	18%	539	13.9–23.7	86	17%	523	12.6–22.0

n = Number of respondents who had 5 or more drinks in one or more occasions in the past 30 days.

% = This is a weighted (adjusted) percentage of the state population (adult) at risk in this demographic subgroup, based on the survey data.

N = Total number of respondents in this subgroup.

95% CI = 95% Confidence Interval; the range of values within which the true value of a prevalence estimate would be expected to fall within, 95% of the time.