

Table 105-0046 [1,2,3,7,8](#)

Males who reported having a prostate-specific antigen (PSA) test, by age group, household population aged 40 and over, Canada, provinces, territories, health regions (January 2000 boundaries) and peer groups, every 2 years

Survey or program details:

Canadian Community Health Survey - [3226](#)

Geography 7,8	Age group	Prostate-specific antigen (PSA) test 2,3	Characteristics 4,5,6,20	2000/2001
Health Labrador Corporation, Newfoundland and Labrador (Peer group F) [1006]	Total, 40 years and over	Total, prostate-specific antigen test	Number of persons	474.0
			Percent	100.0
		Prostate-specific antigen test, less than one year ago	Number of persons	111.3 ^E
			Percent	23.5 ^E
		Prostate-specific antigen test, 1 year to less than 2 years ago	Number of persons	F
			Percent	F
		Prostate-specific antigen test, 2 or more years ago	Number of persons	F
			Percent	F
		Never had a prostate-specific antigen test	Number of persons	274.3
			Percent	57.9
		Prostate-specific antigen test, not stated	Number of persons	F
			Percent	F
	40 to 44 years	Total, prostate-specific antigen test	Number of persons	103.7
			Percent	100.0
		Prostate-specific antigen test, less than one year ago	Number of persons	F
			Percent	F
		Prostate-specific antigen test, 1 year to less than 2 years ago	Number of persons	F
			Percent	F
		Prostate-specific antigen test, 2 or more years ago	Number of persons	F
			Percent	F
		Never had a prostate-specific antigen test	Number of persons	83.8 ^E
			Percent	80.8
		Prostate-specific antigen test, not stated	Number of persons	F
			Percent	F
	45 to 64 years	Total, prostate-specific antigen test	Number of persons	318.2
			Percent	100.0
		Prostate-specific antigen test, less than one year ago	Number of persons	97.8 ^E
			Percent	30.7 ^E
		Prostate-specific antigen test, 1 year to less than 2 years ago	Number of persons	F
			Percent	F
		Prostate-specific antigen test, 2 or more years ago	Number of persons	F
			Percent	F
		Never had a prostate-specific antigen test	Number of persons	158.8
			Percent	49.9
		Prostate-specific antigen test, not stated	Number of persons	F
			Percent	F
		Total, prostate-specific antigen test	Number of persons	F
			Percent	F
		Prostate-specific antigen test, less than one year ago	Number of persons	F

		65 years and over		Percent	F
			Prostate-specific antigen test, 1 year to less than 2 years ago	Number of persons	F
				Percent	F
			Prostate-specific antigen test, 2 or more years ago	Number of persons	F
				Percent	F
			Never had a prostate-specific antigen test	Number of persons	F
				Percent	F
			Prostate-specific antigen test, not stated	Number of persons	F
				Percent	F
Yukon Territory [60]	Total, 40 years and over		Total, prostate-specific antigen test	Number of persons	657.8
				Percent	100.0
			Prostate-specific antigen test, less than one year ago	Number of persons	118.6 ^E
				Percent	18.0 ^E
			Prostate-specific antigen test, 1 year to less than 2 years ago	Number of persons	F
				Percent	F
			Prostate-specific antigen test, 2 or more years ago	Number of persons	39.2 ^E
				Percent	6.0 ^E
			Never had a prostate-specific antigen test	Number of persons	415.7
				Percent	63.2
			Prostate-specific antigen test, not stated	Number of persons	44.1 ^E
				Percent	6.7 ^E
	40 to 44 years		Total, prostate-specific antigen test	Number of persons	175.8
				Percent	100.0
			Prostate-specific antigen test, less than one year ago	Number of persons	F
				Percent	F
			Prostate-specific antigen test, 1 year to less than 2 years ago	Number of persons	F
				Percent	F
			Prostate-specific antigen test, 2 or more years ago	Number of persons	F
				Percent	F
			Never had a prostate-specific antigen test	Number of persons	129.8 ^E
				Percent	73.8
			Prostate-specific antigen test, not stated	Number of persons	F
				Percent	F
	45 to 64 years		Total, prostate-specific antigen test	Number of persons	403.1
				Percent	100.0
			Prostate-specific antigen test, less than one year ago	Number of persons	72.7 ^E
				Percent	18.0 ^E
			Prostate-specific antigen test, 1 year to less than 2 years ago	Number of persons	F
				Percent	F
			Prostate-specific antigen test, 2 or more years ago	Number of persons	F
				Percent	F
			Never had a prostate-specific antigen test	Number of persons	256.5
				Percent	63.6
			Prostate-specific antigen test, not stated	Number of persons	F
				Percent	F
	65 years and over		Total, prostate-specific antigen test	Number of persons	78.9
				Percent	100.0
			Prostate-specific antigen test, less than one year ago	Number of persons	23.2 ^E
				Percent	29.4 ^E
			Prostate-specific antigen test, 1 year to less than 2 years ago	Number of persons	F
				Percent	F
			Prostate-specific antigen test, 2 or more	Number of persons	F

		years ago	Percent	F
		Never had a prostate-specific antigen test	Number of persons	29.3 ^E
			Percent	37.2 ^E
		Prostate-specific antigen test, not stated	Number of persons	F
			Percent	F
Northwest Territories [61]	Total, 40 years and over	Total, prostate-specific antigen test	Number of persons	640.3
			Percent	100.0
		Prostate-specific antigen test, less than one year ago	Number of persons	80.8 ^E
			Percent	12.6 ^E
		Prostate-specific antigen test, 1 year to less than 2 years ago	Number of persons	F
			Percent	F
		Prostate-specific antigen test, 2 or more years ago	Number of persons	F
			Percent	F
		Never had a prostate-specific antigen test	Number of persons	462.0
			Percent	72.1
		Prostate-specific antigen test, not stated	Number of persons	61.1 ^E
			Percent	9.5 ^E
	40 to 44 years	Total, prostate-specific antigen test	Number of persons	156.2
			Percent	100.0
		Prostate-specific antigen test, less than one year ago	Number of persons	F
			Percent	F
		Prostate-specific antigen test, 1 year to less than 2 years ago	Number of persons	F
			Percent	F
		Prostate-specific antigen test, 2 or more years ago	Number of persons	F
			Percent	F
		Never had a prostate-specific antigen test	Number of persons	120.8
			Percent	77.3
		Prostate-specific antigen test, not stated	Number of persons	F
			Percent	F
	45 to 64 years	Total, prostate-specific antigen test	Number of persons	402.1
			Percent	100.0
		Prostate-specific antigen test, less than one year ago	Number of persons	46.9 ^E
			Percent	11.7 ^E
		Prostate-specific antigen test, 1 year to less than 2 years ago	Number of persons	F
			Percent	F
		Prostate-specific antigen test, 2 or more years ago	Number of persons	F
			Percent	F
		Never had a prostate-specific antigen test	Number of persons	292.2
			Percent	72.7
		Prostate-specific antigen test, not stated	Number of persons	F
			Percent	F
	65 years and over	Total, prostate-specific antigen test	Number of persons	82.0
			Percent	100.0
		Prostate-specific antigen test, less than one year ago	Number of persons	F
			Percent	F
		Prostate-specific antigen test, 1 year to less than 2 years ago	Number of persons	F
			Percent	F
		Prostate-specific antigen test, 2 or more years ago	Number of persons	F
			Percent	F
		Never had a prostate-specific antigen test	Number of persons	49.0
			Percent	59.7
		Prostate-specific antigen test, not stated	Number of persons	F

Nunavut [62]	Total, 40 years and over	Total, prostate-specific antigen test	Percent	F
			Number of persons	310.1
		Prostate-specific antigen test, less than one year ago	Percent	100.0
			Number of persons	F
		Prostate-specific antigen test, 1 year to less than 2 years ago	Percent	F
			Number of persons	F
		Prostate-specific antigen test, 2 or more years ago	Percent	F
			Number of persons	F
		Never had a prostate-specific antigen test	Percent	249.7
			Number of persons	80.5
		Prostate-specific antigen test, not stated	Percent	F
			Number of persons	F
	40 to 44 years	Total, prostate-specific antigen test	Percent	75.1
			Number of persons	100.0
		Prostate-specific antigen test, less than one year ago	Percent	F
			Number of persons	F
		Prostate-specific antigen test, 1 year to less than 2 years ago	Percent	F
			Number of persons	F
		Prostate-specific antigen test, 2 or more years ago	Percent	F
			Number of persons	F
		Never had a prostate-specific antigen test	Percent	73.3
			Number of persons	97.6
		Prostate-specific antigen test, not stated	Percent	F
			Number of persons	F
	45 to 64 years	Total, prostate-specific antigen test	Percent	193.7
			Number of persons	100.0
		Prostate-specific antigen test, less than one year ago	Percent	F
			Number of persons	F
		Prostate-specific antigen test, 1 year to less than 2 years ago	Percent	F
			Number of persons	F
		Prostate-specific antigen test, 2 or more years ago	Percent	F
			Number of persons	F
		Never had a prostate-specific antigen test	Percent	150.2
			Number of persons	77.5
		Prostate-specific antigen test, not stated	Percent	F
			Number of persons	F
	65 years and over	Total, prostate-specific antigen test	Percent	41.2
			Number of persons	100.0
		Prostate-specific antigen test, less than one year ago	Percent	F
			Number of persons	F
		Prostate-specific antigen test, 1 year to less than 2 years ago	Percent	F
			Number of persons	F
		Prostate-specific antigen test, 2 or more years ago	Percent	F
			Number of persons	F
		Never had a prostate-specific antigen test	Percent	26.2
			Number of persons	63.6
		Prostate-specific antigen test, not stated	Percent	F
			Number of persons	F

Symbol legend:

^E Use with caution

^F Too unreliable to be published

Footnotes:

1. Source: Statistics Canada, Canadian Community Health Survey, 2000/2001
2. Males aged 40 and over who have had a prostate-specific antigen (PSA) test.
3. A prostate-specific antigen (PSA) test constitutes a blood sample and analysis, not a physical prostate exam.
4. Bootstrapping techniques were used to produce the coefficient of variation (CV) and 95% confidence intervals (CIs).
5. Data with a coefficient of variation (CV) from 16.6% to 33.3% are identified by an (E) and should be interpreted with caution.
6. Data with a coefficient of variation (CV) greater than 33.3% were suppressed (F) due to extreme sampling variability.
7. Health regions are defined by provincial governments as the areas of responsibility for regional health boards (legislated) or as regions of interest to health care authorities.
8. A "peer group" is a grouping of health regions that have similar social and economic characteristics.
20. The following standard symbols are used in this Statistics Canada table: (..) for figures not available for a specific reference period and (...) for figures not applicable.

Source: Statistics Canada. *Table 105-0046 - Males who reported having a prostate-specific antigen (PSA) test, by age group, household population aged 40 and over, Canada, provinces, territories, health regions (January 2000 boundaries) and peer groups, every 2 years*, CANSIM (database).

<http://cansim2.statcan.ca/cgi-win/cnsmcgui.exe?>

[Lang=E&CANSIMFile=CII\CII_1_E.htm&RootDir=CII/](http://cansim2.statcan.ca/cgi-win/cnsmcgui.exe?Lang=E&CANSIMFile=CII\CII_1_E.htm&RootDir=CII/)

(accessed: October 1, 2008)

[Printer-friendly format](#)