

Make another selection | Group by Health Indicator | Download | Printer Friendly

Graph Indicator: Coronary Artery Bypass Graft (CABG) 1997 Graph

Community and Health System Characteristics

[60] Yukon Territory

Indicator	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Coronary Artery Bypass Graft (CABG) - Both sexes Age-standardized rate/100,000 (95% Confidence Interval) Notes: 1 2 3	..	..	..	53.8 (11.5 - 96.1)	96.8 (44.8 - 148.8)	68.8 (27.8 - 109.8)	120.6 (63.0 - 178.2)	100.4 (52.1 - 148.7)	55.1 (22.7 - 87.5)	63.1 (26.6 - 99.5)	62 (26 - 99)	81 (42 - 120)
Coronary Artery Bypass Graft (CABG) - Female Age-standardized rate/100,000 (95% Confidence Interval) Notes: 1 2 3	..	..	..	†	†	†	†	51.2 (9.6 - 92.8)	†	† († - †)	† († - †)	† († - †)
Coronary Artery Bypass Graft (CABG) - Male Age-standardized rate/100,000 (95% Confidence Interval) Notes: 1 2 3	..	..	..	†	134.6 (52.9 - 216.3)	98.7 (34.0 - 163.4)	226.7 (114.3 - 339.1)	141.8 (59.3 - 224.2)	89.7 (31.1 - 148.3)	118.5 (49.5 - 187.5)	94 (28 - 159)	109 (47 - 170)
Hip Replacement - Both sexes Age-standardized rate/100,000 (95% Confidence Interval) Notes: 1 2 4	110.7	70.5	46.2	54.0 (23.3 - 84.8)	60.3 (24.7 - 95.8)	57.3 (21.9 - 92.7)	71.7 (37.6 - 105.8)	57.4 (26.3 - 88.6)	..	..	..	..
Hip Replacement - Female Age-standardized rate/100,000 (95% Confidence Interval) Notes: 1 2 4	168.7	58.7	†	69.6 (15.6 - 123.5)	82.6 (24.8 - 140.5)	80.2 (21.3 - 139.1)	†	59.1 (11.4 - 106.9)	..	..	..	..
Hip Replacement - Male Age-standardized rate/100,000 (95% Confidence Interval) Notes: 1 2 4	62.2	91.3	69.0	40.6 (8.0 - 73.2)	†	†	78.7 (34.1 - 123.3)	53.4 (15.7 - 91.2)	..	..	..	..
Hip Replacement 20+ - Both sexes Age-standardized rate/100,000 (95% Confidence Interval) Notes: 1 2 4 5 6	..	..	..	..	..	..	..	81.6 (36.9 - 126.2)	57.1 ^A (22.3 - 91.9)	109.5 (65.9 - 153.0)	118 (66 - 169)	156 (96 - 216)
Hip Replacement 20+ - Female Age-standardized rate/100,000 (95% Confidence Interval) Notes: 1 2 4 5 6	..	..	..	..	..	..	..	87.0 (17.2 - 156.8)	56.5 (2.3 - 110.6)	133.7 (61.4 - 205.9)	111 (44 - 179)	122 (51 - 192)
Hip Replacement 20+ - Male Age-standardized rate/100,000 (95% Confidence Interval) Notes: 1 2 4 5 6	..	..	..	..	..	..	..	73.5 (21.3 - 125.6)	62.5 (13.0 - 111.9)	87.0 (37.0 - 136.9)	129 (49 - 209)	196 (93 - 299)
Hysterectomy - Female Age-standardized rate/100,000 (95% Confidence Interval) Notes: 1 2 5 6 7 8	..	..	..	592 (446 - 739)	448 (331 - 566)	359 (249 - 469)	458 (332 - 585)	310 (207 - 413)	249 ^A (156 - 342)	320 (221 - 420)	343 (234 - 452)	246 ^A (158 - 333)
Inflow/Outflow Hip Replacement 20+ - Both sexes Ratio Notes: 1 4 5 6	..	..	..	..	..	..	..	..	0.00	0.00	0.00	0.00
Inflow/Outflow PCI - Both sexes Ratio Notes: 1 9	..	..	..	..	..	..	..	..	0.00	0.00	0.00	0.00
Inflow/Outflow Bypass Surgery - Both sexes Ratio Notes: 1	..	..	..	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Inflow/Outflow Hip Replacement - Both sexes Ratio Notes: 1 4	..	..	..	0.00	0.00	0.00	0.00	0.00	..	..	..	..
Inflow/Outflow Hysterectomy - Female Ratio Notes: 1 5 6	..	..	..	1.00	1.03	0.98	0.97	0.95	1.00	0.95	0.98	0.91
Inflow/Outflow Knee Replacement - Both sexes Ratio Notes: 1 4	..	..	..	0.00	0.00	0.00	0.11	0.50	..	..	..	..
Inflow/Outflow Knee Replacement 20+ - Both sexes Ratio Notes: 1 4 5 6	..	..	..	..	..	..	..	..	0.44	0.50	0.43	0.60
Inflow/Outflow Overall - Both sexes Ratio Notes: 1 5 6	..	..	..	0.89	0.87	0.88	0.87	0.89	0.89	0.88	0.84	0.85
Knee Replacement - Both sexes Age-standardized rate/100,000 (95% Confidence Interval) Notes: 1 2 4 10	87.5	48.9	146.6	45.2 (14.2 - 76.2)	77.3 (38.3 - 116.4)	76.8 (37.7 - 115.9)	65.0 (32.2 - 97.8)	44.6 ^A (16.8 - 72.4)	..	..	..	..
Knee Replacement - Female Age-standardized rate/100,000 (95% Confidence Interval) Notes: 1 2 4 10	101.0	59.6	117.5	58.8 (8.2 - 109.4)	82.7 (23.2 - 142.1)	67.6 (16.2 - 119.0)	82.0 (27.9 - 136.1)	49.3 (5.6 - 93.0)	..	..	..	..
Knee Replacement - Male Age-standardized rate/100,000 (95% Confidence Interval) Notes: 1 2 4 10	75.8	†	176.6	†	73.2 (20.9 - 125.6)	83.1 (26.4 - 139.8)	55.0 (10.3 - 99.7)	39.5 (5.0 - 73.9)	..	..	..	..
Knee Replacement 20+ - Both sexes Age-standardized rate/100,000 (95% Confidence Interval)	..	..	..	..	..	..	..	59.8 ^A (22.6 - ..)	103.0 (51.5 - ..)	151.4 (89.8 - ..)	126 (76 - 177)	172 (108 - 235)

[61] Northwest Territories

Indicator	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Coronary Artery Bypass Graft (CABG) - Both sexes Age-standardized rate/100,000 (95% Confidence Interval) Notes: 1 2 3 13	..	..	..	75.1 (29.9 - 120.3)	50.9^ (15.1 - 86.7)	39.6^ (12.8 - 66.3)	97.3 (46.0 - 148.5)	94.9 (45.1 - 144.8)	49.8^ (17.9 - 81.8)	30.6^ (5.7 - 55.6)	88 (43 - 132)	63 (22 - 104)
Coronary Artery Bypass Graft (CABG) - Female Age-standardized rate/100,000 (95% Confidence Interval) Notes: 1 2 3 13	..	..	..	†	†	†	†	†	†	† († - †)	† († - †)	† († - †)
Coronary Artery Bypass Graft (CABG) - Male Age-standardized rate/100,000 (95% Confidence Interval) Notes: 1 2 3 13	..	..	..	103.5 (30.6 - 176.5)	83.5 (20.3 - 146.7)	52.7 (14.3 - 91.0)	165.2 (72.7 - 257.7)	179.3 (84.6 - 274.0)	87.6 (28.8 - 146.4)	58.4^ (11.1 - 105.7)	121 (51 - 191)	115 (35 - 195)
Hip Replacement - Both sexes Age-standardized rate/100,000 (95% Confidence Interval) Notes: 1 2 4 13	29.8	39.7	73.6	37.3 (12.2 - 62.4)	50.6 (22.8 - 78.5)	47.9 (15.1 - 80.7)	97.0 (53.3 - 140.8)	99.1 (58.0 - 140.1)	..	..	..	..
Hip Replacement - Female Age-standardized rate/100,000 (95% Confidence Interval) Notes: 1 2 4 13	36.7	54.5	97.4	42.8 (6.4 - 79.2)	91.0 (37.9 - 144.1)	49.3 (3.8 - 94.8)	122.4 (54.4 - 190.4)	127.9 (60.2 - 195.6)	..	..	..	..
Hip Replacement - Male Age-standardized rate/100,000 (95% Confidence Interval) Notes: 1 2 4 13	†	28.1	52.2	†	†	40.4 (1.0 - 79.8)	77.1 (19.0 - 135.2)	72.6 (24.5 - 120.7)	..	..	..	..
Hip Replacement 20+ - Both sexes Age-standardized rate/100,000 (95% Confidence Interval) Notes: 1 2 4 5 6 13	..	..	..	..	..	..	..	133.6 (78.4 - 188.9)	133.3 (78.3 - 188.2)	71.7 (36.3 - 107.1)	136 (80 - 191)	84 (42 - 126)
Hip Replacement 20+ - Female Age-standardized rate/100,000 (95% Confidence Interval) Notes: 1 2 4 5 6 13	..	..	..	..	..	..	..	180.5 (85.0 - 276.1)	127.8 (48.7 - 206.9)	98.0 (29.1 - 166.9)	169 (77 - 262)	103 (38 - 168)
Hip Replacement 20+ - Male Age-standardized rate/100,000 (95% Confidence Interval) Notes: 1 2 4 5 6 13	..	..	..	..	..	..	..	95.0 (32.7 - 157.3)	141.8 (63.1 - 220.5)	53.7^ (20.3 - 87.1)	103 (41 - 166)	75 (13 - 137)
Hysterectomy - Female Age-standardized rate/100,000 (95% Confidence Interval) Notes: 1 2 5 6 7 8 13	..	..	..	194^ (116 - 273)	247^ (159 - 334)	325 (218 - 431)	210^ (134 - 285)	206^ (130 - 281)	251^ (166 - 335)	491^ (379 - 604)	409 (302 - 517)	371 (267 - 476)
Inflow/Outflow Hip Replacement 20+ - Both sexes Ratio Notes: 1 4 5 6 13	..	..	..	..	..	..	..	..	0.96	0.89	1.00	0.95
Inflow/Outflow PCI - Both sexes Ratio Notes: 1 9 13	..	..	..	..	..	..	..	..	0.00	0.00	0.00	0.00
Inflow/Outflow Bypass Surgery - Both sexes Ratio Notes: 1 13	..	..	..	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Inflow/Outflow Hip Replacement - Both sexes Ratio Notes: 1 4 13	..	..	..	1.10	0.80	0.80	0.55	1.04	..	..	..	..
Inflow/Outflow Hysterectomy - Female Ratio Notes: 1 5 6 13	..	..	..	0.96	0.91	0.88	0.91	0.81	1.03	0.99	0.90	1.08
Inflow/Outflow Knee Replacement - Both sexes Ratio Notes: 1 4 13	..	..	..	1.07	1.21	0.93	1.03	1.34	..	..	..	..
Inflow/Outflow Knee Replacement 20+ - Both sexes Ratio Notes: 1 4 5 6 13	..	..	..	..	..	..	..	..	1.29	1.00	0.93	1.00
Inflow/Outflow Overall - Both sexes	..	..	..	1.05	1.04	0.98	0.97	1.00	0.99	1.00	0.97	0.99

Ratio													
Notes: 1 5 6 13													
Knee Replacement - Both sexes	78.2	107.3	89.3	74.3	93.6	80.5	135.5 ^A	138.8	..	..	..	..	..
Age-standardized rate/100,000 (95% Confidence Interval)				(34.9 - 113.7)	(49.8 - 137.3)	(37.8 - 123.2)	(84.7 - 186.4)	(87.1 - 190.4)					
Notes: 1 2 4 10 13													
Knee Replacement - Female	105.8	94.9	141.1	103.6	104.9	96.9	96.7	160.3	..	..	..	..	..
Age-standardized rate/100,000 (95% Confidence Interval)				(38.7 - 168.4)	(41.5 - 168.3)	(27.7 - 166.1)	(38.1 - 155.2)	(81.1 - 239.5)					
Notes: 1 2 4 10 13													
Knee Replacement - Male	50.3	115.4	†	†	85.8	61.4	177.8	120.7	..	..	..	..	..
Age-standardized rate/100,000 (95% Confidence Interval)					(24.0 - 147.6)	(13.1 - 109.6)	(94.0 - 261.5)	(52.9 - 188.5)					
Notes: 1 2 4 10 13													
Knee Replacement 20+ - Both sexes	..	..	..	..	..	..	..	189.6	156.2	136.7	140	84 ^A	..
Age-standardized rate/100,000 (95% Confidence Interval)								(120.6 - 258.6)	(94.3 - 218.2)	(80.5 - 192.9)	(85 - 194)	(50 - 119)	..
Notes: 1 2 4 5 6 13													
Knee Replacement 20+ - Female	..	..	..	..	..	..	..	231.2	159.1	160.7	155	84 ^A	..
Age-standardized rate/100,000 (95% Confidence Interval)								(121.2 - 341.2)	(68.8 - 249.4)	(74.7 - 246.7)	(73 - 237)	(33 - 135)	..
Notes: 1 2 4 5 6 13													
Knee Replacement 20+ - Male	..	..	..	..	..	..	..	156.9	162.8	121.3	132	82 ^A	..
Age-standardized rate/100,000 (95% Confidence Interval)								(68.8 - 245.0)	(71.8 - 253.8)	(44.8 - 197.9)	(56 - 209)	(37 - 127)	..
Notes: 1 2 4 5 6 13													
Percutaneous Coronary Intervention - Both sexes	..	..	..	..	..	..	192.3	200.2	156.8	121.5	193	144	..
Age-standardized rate/100,000 (95% Confidence Interval)							(126.5 - 258.2)	(134.0 - 266.4)	(100.5 - 213.2)	(74.4 - 168.7)	(130 - 255)	(93 - 195)	..
Notes: 1 2 3 9 13													
Percutaneous Coronary Intervention - Female	..	..	..	..	..	..	100.5	106.7	67.7	55.5	94	98	..
Age-standardized rate/100,000 (95% Confidence Interval)							(26.1 - 174.9)	(34.0 - 179.3)	(8.7 - 126.8)	(10.8 - 100.1)	(22 - 167)	(35 - 160)	..
Notes: 1 2 3 9 13													
Percutaneous Coronary Intervention - Male	..	..	..	..	..	..	273.1	286.2	238.3	181.3	275	185	..
Age-standardized rate/100,000 (95% Confidence Interval)							(167.4 - 378.9)	(177.1 - 395.2)	(143.4 - 333.2)	(100.5 - 262.1)	(178 - 372)	(107 - 264)	..
Notes: 1 2 3 9 13													
General/Family Physicians - Both sexes	..	..	..	71	58	72	69	86	70	83	84	76	..
Rate/100,000 (95% Confidence Interval)				(45 - 97)	(35 - 82)	(47 - 98)	(44 - 94)	(59 - 114)	(45 - 95)	(56 - 111)	(57 - 112)	(50 - 102)	..
Notes: 1 11 13													
Specialist Physicians - Both sexes	..	..	..	44	32	39	33	33	33	31	30	35	..
Rate/100,000 (95% Confidence Interval)				(24 - 64)	(14 - 49)	(20 - 58)	(16 - 51)	(16 - 50)	(16 - 50)	(14 - 48)	(14 - 47)	(17 - 52)	..
Notes: 1 11 13													

[62] Nunavut

Indicator	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Coronary Artery Bypass Graft (CABG) - Both sexes	..	..	..	†	86.2	†	†	30.4 ^A	†	47.1	54	66
Age-standardized rate/100,000 (95% Confidence Interval)					(20.6 - 151.8)			(3.6 - 57.2)		(1.2 - 93.1)	(19 - 152)	(4 - 129)
Notes: 1 2 3 13												
Coronary Artery Bypass Graft (CABG) - Female	..	..	..	†	†	†	†	†	†	†	†	†
Age-standardized rate/100,000 (95% Confidence Interval)										(† - †)	(† - †)	(† - †)
Notes: 1 2 3 13												
Coronary Artery Bypass Graft (CABG) - Male	..	..	..	†	151.7	†	†	62.4	†	89.4	80	†
Age-standardized rate/100,000 (95% Confidence Interval)					(27.5 - 275.8)			(7.4 - 117.4)		(5.1 - 173.7)	(28 - 232)	(† - †)
Notes: 1 2 3 13												
Hip Replacement - Both sexes	29.8	39.7	†	†	54.5	36.0	†	69.0	..	..	..	..
Age-standardized rate/100,000 (95% Confidence Interval)					(12.1 - 97.0)	(2.8 - 69.3)		(5.8 - 132.3)				
Notes: 1 2 4 13												
Hip Replacement - Female	36.7	54.5	†	†	78.9	†	†	†	..	..	..	..
Age-standardized rate/100,000 (95% Confidence Interval)					(2.1 - 155.8)							
Notes: 1 2 4 13												
Hip Replacement - Male	†	28.1	†	†	†	†	†	†	..	..	..	..
Age-standardized rate/100,000 (95% Confidence Interval)												
Notes: 1 2 4 13												
Hip Replacement 20+ - Both sexes	..	..	..	..	..	..	..	92.9	†	118.7	†	59
Age-standardized rate/100,000 (95% Confidence Interval)								(8.1 - 177.7)		(35.5 - 201.9)	(† - †)	(15 - 103)
Notes: 1 2 4 5 6 13												
Hip Replacement 20+ - Female	..	..	..	..	..	..	..	†	†	170.5	†	†
Age-standardized rate/100,000 (95% Confidence Interval)										(55.6 - 521.6)	(† - †)	(† - †)
Notes: 1 2 4 5 6 13												
Hip Replacement 20+ - Male	..	..	..	..	..	..	..	†	†	97.7	†	78
Age-standardized rate/100,000 (95% Confidence Interval)										(8.7 - 186.7)	(† - †)	(5 - 151)
Notes: 1 2 4 5 6 13												
Hysterectomy - Female	..	..	..	225 ^A	231 ^A	..	168 ^A	231 ^A	274	193 ^A	172 ^A	185 ^A
Age-standardized rate/100,000 (95% Confidence Interval)				(90 - 360)	(108 - 353)		(40 - 295)	(119 - 342)	(159 - 389)	(101 - 285)	(85 - 260)	(72 - 299)
Notes: 1 2 5 6 7 8 13												
Inflow/Outflow Hip Replacement 20+ - Both sexes	..	..	..	..	..	..	..	..	0.00	0.00	0.00	0.00
Ratio												
Notes: 1 4 5 6 13												
Inflow/Outflow PCI - Both sexes	..	..	..	..	..	..	..	..	0.00	0.00	0.00	0.00
Ratio												
Notes: 1 9 13												
Inflow/Outflow Bypass Surgery - Both sexes	..	..	..	0.00	0.00	..	0.00	0.00	0.00	0.00	0.00	0.00
Ratio												
Notes: 1 13												
Inflow/Outflow Hip Replacement - Both sexes	..	..	..	0.00	0.00	..	0.00	0.00	0.00	0.00	0.00	0.00

Inflow/Outflow Hip Replacement - Both sexes	..	..	..	0.00	0.00	..	0.00	0.00	..	..	..	..
Ratio												
Notes: 1 4 13												
Inflow/Outflow Hysterectomy - Female	..	..	..	0.69	0.33	..	0.22	0.58	0.52	0.50	0.20	0.29
Ratio												
Notes: 1 5 6 13												
Inflow/Outflow Knee Replacement - Both sexes	..	..	..	0.00	0.00	..	0.00	0.00	..	..	..	..
Ratio												
Notes: 1 4 13												
Inflow/Outflow Knee Replacement 20+ - Both sexes	..	..	..	..	..	..	..	..	0.00	0.00	0.00	0.00
Ratio												
Notes: 1 4 5 6 13												
Inflow/Outflow Overall - Both sexes	..	..	..	0.59	0.49	..	0.51	0.41	0.43	0.43	0.43	0.46
Ratio												
Notes: 1 5 6 13												
Knee Replacement - Both sexes	78.2	107.3	83.7	76.0 (17.6 - 134.5)	153.9 [^] (71.8 - 236.1)	84.7 (31.9 - 137.6)	117.2 (31.2 - 203.3)	278.6 [^] (173.7 - 383.5)	..	..	..	..
Age-standardized rate/100,000 (95% Confidence Interval)												
Notes: 1 2 4 10 13												
Knee Replacement - Female	105.8	94.9	131.9	111.6 (19.5 - 203.8)	240.9 (72.3 - 409.6)	182.8 (60.1 - 305.5)	†	482.2 (250.9 - 713.5)	..	..	..	..
Age-standardized rate/100,000 (95% Confidence Interval)												
Notes: 1 2 4 10 13												
Knee Replacement - Male	50.3	115.4	†	†	81.9 (18.7 - 145.1)	†	†	142.1 (48.6 - 235.6)	..	..	..	..
Age-standardized rate/100,000 (95% Confidence Interval)												
Notes: 1 2 4 10 13												
Knee Replacement 20+ - Both sexes	..	..	..	..	..	..	..	343.1 [^] (209.1 - 477.0)	238.8 (137.0 - 340.7)	418.0 [^] (263.7 - 572.3)	394 [^] (232 - 557)	358 [^] (227 - 489)
Age-standardized rate/100,000 (95% Confidence Interval)												
Notes: 1 2 4 5 6 13												
Knee Replacement 20+ - Female	..	..	..	..	..	..	..	544.0 [^] (272.3 - 815.7)	267.0 (109.4 - 424.6)	737.4 [^] (416.5 - 1058.2)	692 [^] (351 - 1032)	491 [^] (261 - 720)
Age-standardized rate/100,000 (95% Confidence Interval)												
Notes: 1 2 4 5 6 13												
Knee Replacement 20+ - Male	..	..	..	..	..	..	..	191.7 (64.0 - 319.3)	214.0 (86.2 - 341.8)	170.5 (35.8 - 305.3)	165 (32 - 299)	241 (97 - 385)
Age-standardized rate/100,000 (95% Confidence Interval)												
Notes: 1 2 4 5 6 13												
Percutaneous Coronary Intervention - Both sexes	..	..	..	..	..	..	100.8 (31.2 - 170.4)	120.2 (42.0 - 198.5)	85.8 [^] (37.9 - 133.7)	68.1 (24.7 - 111.5)	109 (48 - 171)	105 (41 - 168)
Age-standardized rate/100,000 (95% Confidence Interval)												
Notes: 1 2 3 9 13												
Percutaneous Coronary Intervention - Female	..	..	..	..	..	..	†	†	†	†	†	†
Age-standardized rate/100,000 (95% Confidence Interval)												
Notes: 1 2 3 9 13												
Percutaneous Coronary Intervention - Male	..	..	..	..	..	..	154.9 (50.8 - 259.0)	208.2 (80.0 - 336.4)	147.0 [^] (61.4 - 232.7)	114.1 (36.5 - 191.6)	131 (49 - 212)	79 [^] (9 - 149)
Age-standardized rate/100,000 (95% Confidence Interval)												
Notes: 1 2 3 9 13												
General/Family Physicians - Both sexes	..	..	..	22 (4 - 39)	25 (6 - 43)	35 (13 - 56)	34 (13 - 56)	24 (6 - 41)	43 (20 - 67)	32 (12 - 53)	26 (8 - 44)	32 (12 - 52)
Rate/100,000 (95% Confidence Interval)												
Notes: 1 11 13												
Specialist Physicians - Both sexes	..	..	..	†	0 (0 - 0)	†	†	†	†	†	†	†
Rate/100,000 (95% Confidence Interval)												
Notes: 1 11 13												

Back to Search

Close

PLEASE NOTE:

For further information including maps, definitions, and data sources please click on region name and indicator name links.

All indicators with the exception of those specific to physicians are based on a data collection period of April 1 - March 31. Physician indicators derive from the period January 1 - December 31.

† Figures suppressed due to small numbers or incomplete data

.. Figures not available

[^] Statistically different from the average (Canada) rate ($p \leq 0.05$)

‡ The Canada rate includes only those provinces/territories for which comparable data were available.

* Statistically significant Disparity rate ratio or Potential rate reduction

1. For PEI, the Yukon, the Northwest Territories and Nunavut, the entire province or territory is included as a region.

2. Canadian, provincial and territorial rates are based on October 1 population estimates. Regional rates are based on July 1 population estimates.

3. Cardiac procedures for residents of Zone 4 and 5, New Brunswick, are performed mostly in Quebec; rates are suppressed due to differences in data collection and unavailability of data from Quebec.

4. Beginning with data year 2004, the indicator was calculated for the population aged 20 years and older and therefore is not comparable with the previous rates for the indicator which included all ages.

5. Hospitalization data for 2006–2007 for Peace Country Health Region in Alberta were incomplete therefore the indicator could not be calculated for this region.

6. Hospitalization data for 2008–2009 for Area 8 (Peace Country) in Alberta was incomplete, therefore the indicator could not be calculated for this region.

7. Hysterectomy rates for 2001 to 2005 may not be comparable to other years due to a change in the methodology.

8. Beginning with data year 2006–2007, hysterectomy rates include both total and sub-total hysterectomies, similar to the reporting prior to 2001–2002 data. Comparison with rates reported for previous years should be made with caution.

9. Rates for Quebec are not available due to differences in data collection.

10. Beginning with data year 2003, rates include revisions to knee replacements for all provinces/territories. Previously, rates for Quebec may have been underestimated due to data issues, which did not permit the identification of knee replacement revisions.

11. Canadian, provincial and regional rates are based on July 1 population estimates.

12. Due to licensing authority data supply interruptions in 2000, rates for Alberta and the Yukon may be underestimated.

13. For the Northwest Territories and Nunavut, values derived from 1997 or 1998 data years are based on the combined data from these jurisdictions.