

Suicide

Year	Geography	Measure	Age	Gender	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	
Health Region North	number of deaths	all ages	men	21	20	17	24	23	24	25	34	18	26	46	47	50	50	55	49	47	63	53	55	53		
			women	10	3	7	7	6	13	8	7	17	9	8	13	14	10	16	12	10	17	16	17			
		0-14 yrs	men	0	0	0	0	0	1	0	0	0	0	0	0	0	1	4	2	2	0	1	0	1	0	1
			women	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	2	1	0
		15-24 yrs	men	1	4	5	2	1	4	3	5	3	3	3	6	16	6	10	11	11	7	21	19	13	7	
			women	0	1	2	0	1	4	3	1	1	3	1	0	2	1	0	3	0	2	2	2	2	3	
		25-44 yrs	men	12	5	1	5	11	8	5	13	5	7	10	11	15	21	24	24	25	19	17	17	18		
			women	3	2	3	2	2	3	2	4	2	6	2	5	6	3	3	6	3	1	5	3	4		
		45-64 yrs	men	6	8	7	10	9	6	11	10	6	13	22	15	17	11	15	11	5	15	9	19	17		
			women	5	0	1	5	2	6	2	1	4	7	4	2	3	5	4	6	3	3	5	4	5		
		65+ yrs	men	2	3	4	7	2	5	6	6	4	3	8	4	8	6	3	3	9	8	7	6	10		
			women	2	0	1	0	1	0	1	2	0	1	2	1	2	5	2	1	6	3	3	6	5		
		deaths per 100 000 persons	all ages	men	10	9	8	12	11	11	12	15	8	12	20	20	22	21	23	20	19	26	22	23	23	
				women	5	1	4	3	3	6	4	4	3	8	4	4	6	6	4	7	5	4	8	7	7	
	0-14 yrs		men	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
			women	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	15-24 yrs		men	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
			women	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	25-44 yrs		men	22	8	2	9	21	13	8	22	8	11	15	16	22	30	34	36	36	26	23	23	25		
			women	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	45-64 yrs		men	12	16	14	21	17	12	22	20	12	28	46	31	36	24	33	25	10	35	20	42	39		
			women	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	65+ yrs		men	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
			women	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		

Year	Geography	Measure	Age	Gender	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Health Region North	number of deaths	all ages	men	53	50	63	54	40	54	40	47	40	40	49	37	37	44	33	36	26	39	45	36	
			women	20	14	10	15	13	12	14	16	13	11	10	10	14	11	22	13	3	15	12	13	
		0-14 yrs	men	0	0	2	2	1	2	0	0	0	0	0	0	0	0	0	2	0	1	0	0	0
			women	0	1	0	1	0	1	0	1	0	0	0	0	1	0	0	0	0	0	0	0	0
		15-24 yrs	men	13	11	15	13	8	8	9	14	6	8	7	3	5	9	5	7	2	7	5	7	
			women	3	1	3	2	5	1	3	1	3	2	5	3	3	3	4	2	1	5	2	1	
		25-44 yrs	men	18	15	27	18	16	23	9	16	13	15	18	15	20	20	16	13	8	17	18	11	
			women	7	3	3	5	3	3	6	6	4	5	3	4	5	1	8	5	0	5	5	6	
		45-64 yrs	men	12	15	10	10	11	12	14	8	15	10	17	13	11	11	9	11	13	10	13	12	
			women	6	5	4	3	4	6	5	6	2	1	2	3	5	5	7	4	2	5	5	5	
		65+ yrs	men	10	9	9	11	4	9	8	9	6	7	7	6	1	4	1	5	2	5	9	6	
			women	4	4	0	4	1	1	0	3	4	3	0	0	0	2	3	2	0	0	0	1	
		deaths per 100 000 persons	all ages	men	22	20	27	23	17	22	19	21	17	17	20	16	15	18	14	15	10	16	19	15
				women	9	6	5	6	5	6	6	6	5	5	5	6	4	10	6	1	7	5	5	
	0-14 yrs		men	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
			women	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	15-24 yrs		men	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
			women	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	25-44 yrs		men	26	21	38	25	21	31	12	24	19	22	25	23	30	29	27	21	14	26	29	18	
			women	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	45-64 yrs		men	25	29	21	21	23	22	33	15	27	18	29	22	19	18	14	18	19	16	21	19	
			women	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	65+ yrs		men	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
			women	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

Description

Suicide and intentional self-harm (ICD-10: X60-X84). Number of deaths and deaths per 100 000 persons of same age and gender per year.

The statistics include all persons registered as a resident of Norway at the time of death, regardless of whether the death occurred in Norway or abroad. Rates are age standardised to reduce the effect of differing age distributions when groups are compared over time and between geographical regions. Annual figures are standardised

according to 5-year age groups in the Norwegian population per 1 January 1981.

To view deaths per 100 000 persons, click on the Measures menu, select 'filter' and then 'deaths per 100 000'.

Notes

The entire time series from 1970 is removed if over half the rates are based on fewer than 11 deaths. This is because of reduced reliability.

The following editions of the ICD system have been used: 1970-1985: ICD-8, 1986-1995: ICD-9, 1996 onwards: ICD-10.

Source

Cause of Death Registry, Norwegian Institute of Public Health

Collection

Based on information from the Cause of Death Registry. The Norwegian Institute of Public Health is responsible for the registry, while Statistics Norway performs the data processing.

Mortality statistics are collated from death reports completed by public physicians. Additional information is gathered from results of autopsies and forensic examinations. Cause of death is coded according to the underlying cause of death cited by the physician who signs the death certificate. Underlying causes of death are defined as the illness or injury that, in the opinion of the physician, started the chain of conditions leading directly to death. Causes of death are coded according to the principles and directions of the World Health Organisation (WHO). The international classification system ICD is used.

Data quality

Good.

Time periods

1970-2010

Geographical level

Country, health region

Gender

Men, women

Age groups

Age at death: All ages, 0-14, 15-24, 25-44, 45-64, 65 years and over

Frequency of updates

Annually

Last updated

#####

Keywords

Click on a keyword to search for similar indicators.

[Cause of death](#)

[Dead](#)

[Death](#)

[Diagnosis](#)

[External cause](#)

[Suicide](#)

[Violent death](#)

[Voldsomme dødsfall](#)

[Ytre årsaker](#)