## **Chronic Disease Indicators: State/Area Profile**



**Alaska compared with United States** 

			United
Indicator	Measure 0	Alaska	States
Fruit and vegetable consumption among adults aged >=18 years - 2009 view definition	Prevalence (CI)	<b>23.4%</b> (20.7-26.1)	23.4% <sup>a</sup>
Fruit and vegetable consumption among youth - 2009 <u>view definition</u>	Prevalence (CI)	<b>17.2%</b> (15.1-19.5)	<b>22.3%</b> (21.1-23.7)
Obesity among adults aged >= 18 years - 2010 view definition	Prevalence (CI)	<b>25.2%</b> (22.2-28.2)	27.5% <sup>a</sup>
Overweight or obesity among adults aged >= 18 years - 2010 view definition	Prevalence (CI)	<b>65.9%</b> (62.2-69.3)	64.5% <sup>a</sup>
Obesity among youth - 2009 <u>view definition</u>	Prevalence (CI)	<b>11.8%</b> (9.9-13.9)	<b>12.0%</b> (10.9-13.1)
Recommended physical activity among adults aged >=18 years - 2009 view definition	Prevalence (CI)	<b>60.7%</b> (57.7-63.7)	51.0% <sup>a</sup>
Recommended physical activity among youth - 2009 view definition	Prevalence (CI)	<b>20.2%</b> (17.4-23.3)	<b>18.4%</b> (17.3-19.5)
Television viewing among youth - 2009 <u>view definition</u>	Prevalence (CI)	<b>24.8%</b> (22.8-26.9)	<b>32.8%</b> (30.4-35.3)
Tobacco and Alcohol			
Indicator	Measure <b>1</b>	Alaska	United States
Alcohol use among youth - 2009 <u>view definition</u>	Prevalence (CI)	<b>33.2%</b> (30.0-36.6)	<b>41.8%</b> (40.2-43.4)
Binge drinking among youth - 2009 <u>view definition</u>	Prevalence (CI)	<b>21.7%</b> (18.8-24.8)	<b>24.2%</b> (22.6-25.9)
Binge drinking among adults aged >= 18 years - 2010 view definition	Prevalence (CI)	<b>20.6%</b> (17.4-23.7)	15.1% <sup>a</sup>
Binge drinking among women of childbearing age - 2010 view definition	Prevalence (CI)	<b>18.6%</b> (13.6-24.9)	15.4% <sup>a</sup>
Heavy drinking among adult females aged >= 18 years - 2010 view definition	Prevalence (CI)	<b>4.0%</b> (2.6-6.2)	4.5% <sup>a</sup>
Heavy drinking among adult males aged >= 18 years - 2010 view definition	Prevalence (CI)	<b>7.1%</b> (4.7-10.7)	5.4% <sup>a</sup>
	Number	70	29,165
Chronic liver disease, mortality - 2007 view definition	Crude Rate (CI)	<b>10.2</b> (8.0-12.9)	<b>9.7</b> (9.6-9.8)
	Age-adjusted Rate (CI)	<b>11.4</b> (1.1-1.8)	<b>9.1</b> (9.0-9.2)
Cigarette smoking among adults aged >= 18 years - 2010 view definition	Prevalence (CI)	<b>20.4%</b> (17.3-23.4)	17.3% <sup>a</sup>
Cigarette smoking among youth - 2009 <u>view definition</u>	Prevalence (CI)	<b>15.7%</b> (13.1-18.8)	<b>19.5%</b> (17.9-21.2)
Smokeless tobacco use among youth - 2009 <u>view definition</u>	Prevalence (CI)	<b>13.6%</b> (10.7-17.1)	<b>8.9%</b> (7.3-10.8)
Sale of cigarette packs - 2011 <u>view definition</u>	Packs Sold per Capita	43.3	-
I			
Cancer			
Indicator	Measure	Alaska	United States
	Number	2,440	-
Invasive cancer (all sites combined), incidence - 2004-2008 view definition	Crude Rate (CI)	<b>361.0</b> (354.6-367.5)	<b>481.7</b> (481.4-482.1
	Age-adjusted Rate (CI)	<b>475.6</b> (466.5-484.9)	<b>465.0</b> (464.7-465.4
	Number	791	560,278
Cancer (all sites combined), mortality - 2004-2008 view definition	Crude Rate (CI)	<b>117.1</b> (113.5-120.8)	<b>187.6</b> (187.4-187.8
	Age-adjusted Rate	181.2	181.3

	(CI)		(181.1-181.5)
	Number Crude Rate	105 15.5	21.8
ancer of the bladder (in situ and invasive), incidence - 2004-2008 view definition	(CI)	(14.2-16.9) <b>23.0</b>	(21.8-21.9) <b>21.2</b>
	(CI)	(21.0-25.2)	(21.1-21.3)
	Number Crude Rate	15 2.2	13,527 4.5
ancer of the bladder, mortality - 2004-2008 view definition	(CI)	(1.8-2.8)	(4.5-4.6)
	Age-adjusted Rate (CI)	<b>4.0</b> (3.1-5.0)	<b>4.4</b> (4.3-4.4)
	Number	371	-
nvasive cancer of the female breast, incidence - 2004-2008 view definition	Crude Rate (CI)	<b>114.3</b> (109.2-119.6)	<b>133.3</b> (133.1-133.6)
	Age-adjusted Rate (CI)	<b>130.4</b> (124.2-136.9)	<b>121.0</b> (120.8-121.3)
	Number	56	40,815
ancer of the female breast, mortality - 2004-2008 view definition	Crude Rate (CI)	<b>17.4</b> (15.4-19.6)	<b>26.9</b> (26.8-27.0)
	Age-adjusted Rate (CI)	<b>21.7</b> (19.1-24.5)	<b>23.5</b> (23.4-23.6)
	Number	26	-
nvasive cancer of the cervix, incidence - 2004-2008 view definition	Crude Rate (CI)	<b>8.0</b> (6.7-9.5)	<b>8.3</b> (8.2-8.3)
	Age-adjusted Rate	<b>8.3</b> (6.9-9.9)	<b>8.1</b> (8.0-8.1)
	Number	6	3,956
Cancer of the cervix, mortality - 2004-2008 view definition	Crude Rate (CI)	<b>2.0</b> (1.4-2.8)	<b>2.6</b> (2.6-2.6)
	Age-adjusted Rate	<b>2.2</b> (1.5-3.2)	<b>2.4</b> (2.4-2.4)
	Number	237	-
ancer of the colon and rectum, incidence - 2004-2008 view definition	Crude Rate (CI)	<b>35.1</b> (33.1-37.2)	<b>49.3</b> (49.2-49.5)
· · · · · · · · · · · · · · · · · · ·	Age-adjusted Rate	<b>50.2</b> (47.2-53.4)	<b>47.6</b> (47.5-47.7)
	Number	70	53,171
ancer of the colon and rectum, mortality - 2004-2008 view definition	Crude Rate (CI)	<b>10.3</b> (9.2-11.4)	<b>17.8</b> (17.7-17.9)
	Age-adjusted Rate (CI)	<b>17.2</b> (15.4-19.3)	<b>17.1</b> (17.1-17.2)
	Number	335	-
ancer of the lung and bronchus, incidence - 2004-2008 view definition	Crude Rate (CI)	<b>49.6</b> (47.3-52.0)	<b>69.9</b> (69.8-70.0)
	Age-adjusted Rate	<b>74.0</b> (70.3-77.9)	<b>67.9</b> (67.7-68.0)
	Number	232	158,619
Cancer of the lung and bronchus, mortality - 2004-2008 view definition	Crude Rate (CI)	<b>34.4</b> (32.4-36.4)	<b>53.1</b> (53.0-53.2)
	Age-adjusted Rate (CI)	<b>53.5</b> (50.3-56.9)	<b>51.6</b> (51.5-51.7)
	Number	62	-
nvasive melanoma, incidence - 2004-2008 <u>view definition</u>	Crude Rate (CI)	<b>9.2</b> (8.2-10.3)	<b>19.3</b> (19.2-19.4)
view definition	Age-adjusted Rate	<b>10.6</b> (9.4-12.0)	<b>18.8</b> (18.7-18.8)
	Number	(9.4-12.0)	8,364
Melanoma, mortality - 2004-2008 <u>view definition</u>	Crude Rate (CI)	<b>1.6</b> (1.2-2.1)	<b>2.8</b> (2.8-2.8)
	Age-adjusted Rate	<b>2.1</b> (1.6-2.8)	<b>2.7</b> (2.7-2.7)
	Number	60	-
nvasive cancer of the oral cavity or pharynx, incidence - 2004-2008 view definition	Crude Rate (CI)	<b>8.8</b> (7.9-9.9)	<b>11.3</b> (11.3-11.4)
	, ,	· ,	· ,

	(CI)	(9.4-12.1)	(10.7-10.8)
	Number	11	7,881
Cancer of the oral cavity and pharynx, mortality - 2004-2008 view definition	Crude Rate (CI)	<b>1.6</b> (1.2-2.1)	<b>2.6</b> (2.6-2.7)
	Age-adjusted Rate	<b>2.4</b> (1.8-3.1)	<b>2.5</b> (2.5-2.5)
	Number	352	-
Invasive cancer of the prostate, incidence - 2004-2008 view definition	Crude Rate (CI)	<b>100.1</b> (95.5-104.9)	<b>144.9</b> (144.6-145.1)
200 - 200 -	Age-adjusted Rate (CI)	141.5	<b>152.7</b> (152.4-153.0)
	Number	32	28,769
Cancer of the prostate, mortality - 2004-2008 view definition	Crude Rate (CI)	<b>9.2</b> (7.8-10.7)	<b>19.6</b> (19.5-19.7)
	Age-adjusted Rate (CI)	<b>22.5</b> (18.9-26.5)	<b>24.4</b> (24.3-24.5)
Clinical breast examination among women >= 40 years - 2010 <u>view definition</u>	Prevalence (CI)	<b>79.1%</b> (74.8-82.9)	76.5% <sup>a</sup>
Mammography use among women aged >= 40 years - 2010 view definition	Prevalence (CI)	<b>72.4%</b> (67.7-76.7)	75.6% <sup>a</sup>
Papanicolaou smear use among adult women aged >= 18 years - 2010 view definition	Prevalence (CI)	<b>81.1%</b> (76.0-85.3)	81.1% <sup>a</sup>
Fecal occult blood test among adults aged >= 50 years - 2010 view definition	Prevalence (CI)	<b>7.2%</b> (5.0-10.2)	11.0% <sup>a</sup>
Fecal occult blood test or sigmoidoscopy/colonoscopy among adults aged >= 50 years - 2010 view definition	Prevalence (CI)	<b>49.8%</b> (45.3-54.3)	57.8% <sup>a</sup>
Sigmoidoscopy/colonoscopy among adults >= 50 years - 2010 <u>view definition</u>	Prevalence (CI)	<b>48.0%</b> (43.5-52.5)	52.8% <sup>a</sup>
1			
Cardiovascular Disease			United
Indicator	Measure	Alaska	States 1
	Number	157	135,952
Cerebrovascular disease (Stroke), mortality - 2007 <u>view definition</u>	Crude Rate (CI)	<b>23.0</b> (19.4-26.6)	<b>45.1</b> (44.8-45.3)
	Age-adjusted Rate (CI)	<b>44.3</b> (37.3-51.2)	<b>42.2</b> (42.0-42.5)
	Number	291	277,193
Congestive heart failure, mortality - 2007 <u>view definition</u>	Crude Rate (CI)	<b>42.6</b> (37.7-47.5)	<b>91.9</b> (91.6-92.2)
	Age-adjusted Rate (CI)	<b>85.6</b> (75.8-95.5)	<b>85.4</b> (85.1-85.7)
	Number	406	437,131
Coronary heart disease, mortality - 2007 <u>view definition</u>	Crude Rate (CI)	<b>59.4</b> (53.6-65.2)	<b>144.9</b> (144.5-145.4)
	Age-adjusted Rate (CI)	<b>94.3</b> (85.1-103.5)	<b>135.5</b> (135.1-135.9)
	Number	613	616,067
Diseases of the heart, mortality - 2007 <u>view definition</u>	Crude Rate (CI)	<b>89.7</b> (82.6-96.8)	<b>204.3</b> (203.7-204.8)
	Age-adjusted Rate (CI)	<b>147.9</b> (136.2-159.6)	<b>190.9</b> (190.4-191.4)
	Number	816	806,156
Major cardiovascular diseases, mortality - 2007 <u>view definition</u>	Crude Rate (CI)	<b>119.4</b> (111.2-127.6)	<b>267.3</b> (266.7-267.9)
	Age-adjusted Rate (CI)	<b>203.5</b> (189.5-217.4)	<b>249.9</b> (249.4-250.5)
	Number	475	387,633
Hospitalization for cerebrovascular accident or stroke among Medicare-eligible persons aged >= 65 years - 2006 view definition	Crude Rate (CI)	<b>11.0</b> (10.0-12.0)	<b>13.7</b> (13.6-13.7)
	Age-adjusted Rate (CI)	<b>11.6</b> (10.6-12.7)	<b>13.6</b> (13.5-13.6)
	Number	513	548,313
	Crude Rate	11.9	19.3

- 2006 <u>view definition</u>	Age-adjusted Rate (CI)	<b>13.1</b> (12.0-14.3)	<b>19.2</b> (19.1-19.2)
high blood pressure awareness among adults aged >= 18 years - 2009 view definition	Prevalence (CI)	<b>26.4%</b> (23.9-28.9)	28.7% <sup>a</sup>
Taking medicine for high blood pressure control among adults aged >= 18 years - 2009 view definition	Prevalence (CI)	<b>66.9%</b> (62.0-71.8)	79.2% <sup>a</sup>
Cholesterol screening among adults aged >= 18 years - 2009 <u>view definition</u>	Prevalence (CI)	<b>71.3%</b> (68.1-74.5)	77.0% <sup>a</sup>
Diabetes			
indicator	Measure	Alaska	United States
Diabetes prevalence among adults aged >= 18 years - 2010 <u>view definition</u>	Prevalence (CI)	<b>5.3%</b> (3.9-6.6)	8.7% <sup>a</sup>
	Number	260	231,402
Mortality with diabetes - 2007 <u>view definition</u>	Crude Rate (CI)	<b>38.0</b> (33.4-42.7)	<b>76.7</b> (76.4-77.0)
	Age-adjusted Rate (CI)	<b>61.4</b> (53.9-68.9)	<b>73.1</b> (72.8-73.4)
	Prevalence (CI)	<b>58.1%</b> (49.7-66.5)	-
Dilated eye examination among adults aged >= 18 years with diabetes - 2010 view definition	Age-adjusted Prevalence (CI)	<b>58.3%</b> (49.3-67.3)	<b>62.8%</b> (60.6-65.0)
	Prevalence (CI)	<b>71.8%</b> (64.0-79.6)	-
Foot examination among adults aged >= 18 years with diabetes - 2010 view definition	Age-adjusted Prevalence (CI)	<b>71.3%</b> (62.7-79.9)	<b>67.5%</b> (65.3-69.7)
	Prevalence (CI)	<b>65.2%</b> (57.9-72.5)	-
nfluenza vaccination among adults aged >= 18 years with diabetes - 2010 view definition	Age-adjusted Prevalence (CI)	<b>62.0%</b> (53.6-70.4)	<b>50.1%</b> (48.5-51.7)
Pneumococcal vaccination among adults aged >= 18 years with diabetes - 2010	Prevalence (CI)	<b>63.0%</b> (54.8-71.2)	-
view definition	Age-adjusted Prevalence (CI)	<b>58.4%</b> (49.2-67.6)	-
	Prevalence (CI)	<b>67.3%</b> (59.5-75.1)	-
Self blood glucose monitoring among adults aged >= 18 years with diabetes - 2010 view definition	Age-adjusted Prevalence (CI)	<b>65.9%</b> (57.3-74.5)	<b>63.6%</b> (61.4-65.8)
			ı
Arthritis			
indicator	Measure	Alaska	United States
Arthritis among adults aged >= 18 years - 2009 <u>view definition</u>	Prevalence (CI)	<b>23.5%</b> (20.9-26.0)	<b>25.9%</b> (25.7-26.2)
Activity limitation due to arthritis among adults aged >=18 years - 2009 view definition	Prevalence (CI)	<b>11.6%</b> (9.9-13.2)	<b>11.3%</b> (11.1-11.4)
Physical inactivity among adults aged >=18 years with arthritis - 2009 <u>view definition</u>	Prevalence (CI)	<b>13.8%</b> (10.6-17.7)	<b>19.0%</b> (18.6-19.4)
Arthritis among adults aged >=18 years who are obese - 2009 view definition	Prevalence (CI)	<b>36.6%</b> (31.4-41.7)	<b>34.4%</b> (33.9-34.9)
Arthritis among adults aged >=18 years who have diabetes - 2009 view definition	Prevalence (CI)	<b>39.6%</b> (29.7-49.6)	<b>48.6%</b> (47.7-49.5)
Arthritis among adults aged >=18 years who have heart disease - 2009 view definition	Prevalence (CI)	<b>64.6%</b> (53.5-75.8)	<b>56.5%</b> (55.5-57.5)
air or poor health among adults aged >=18 years with arthritis - 2009 view definition	Prevalence (CI)	<b>25.0%</b> (20.7-30.0)	<b>30.2%</b> (29.8-30.7)
Adults aged >=18 years with arthritis who have taken a class to learn how to manage arthritis symptoms - 2009 view definition	Prevalence (CI)	-	<b>13.0%</b> <sup>b</sup> (12.2-13.8)
Overarching Conditions			

Indicator	Measure	Alaska	States
Current lack of health insurance among adults aged 18-64 years - 2010 view definition	Prevalence (CI)	<b>19.5%</b> (16.2-22.8)	17.8% <sup>a</sup>
High school completion among adults 18-24 years - 2010 view definition	Percent (CI)	81.2%	83.2%
Poverty - 2010 <u>view definition</u>	Percent (CI)	9.9%	15.3%
Life expectancy at birth - 1999-2001 view definition	Years	76.6	77.3
Life expectancy at age 65 years - 1999-2001 view definition	Years	18.0	18.1
	Number	1,036	471,796
Premature mortality among adults aged 45-64 years - 2007 view definition	Crude Rate (CI)	<b>569.9</b> (535.2-604.6)	<b>616.0</b> (614.3-617.8)
	Age-adjusted Rate (CI)	<b>569.5</b> (534.9-604.2)	<b>600.4</b> (598.7-602.1)
	Prevalence (CI)	<b>11.1%</b> (9.2-13.2)	<b>16.1%</b> (15.9-16.4)
Fair or poor self-rated health status among adults aged >= 18 years - 2010 view definition	Age-adjusted Prevalence (CI)	<b>11.7%</b> (9.8-13.8)	<b>15.4%</b> (15.1-15.6)
	Mean Days (CI)	<b>3.2</b> (2.6-3.7)	<b>3.7</b> (3.6-3.8)
Recent physically unhealthy days among adults aged >=18 years - 2010 view definition	Age-adjusted Mean Days (CI)	<b>3.4</b> (2.9-4.0)	<b>3.5</b> (3.4-3.6)
	Mean Days (CI)	<b>2.9</b> (2.4-3.5)	<b>3.5</b> (3.4-3.6)
Recent mentally unhealthy days among adults aged >=18 years - 2010 view definition	Age-adjusted Mean Days (CI)	<b>2.9</b> (2.4-3.4)	<b>3.6</b> (3.5-3.7)
	Mean Days (CI)	<b>2.0</b> (1.6-2.4)	<b>2.3</b> (2.2-2.4)
Recent activity limitation among adults aged >=18 years - 2010 view definition	Age-adjusted Mean Days (CI)	<b>2.1</b> (1.7-2.6)	<b>2.2</b> (2.2-2.3)
Other Diseases and Risk Factors			
Indicator	Measure	Alaska	United States

Indicator	Measure	Alaska	United States
	Number	9	3,447
Asthma, mortality - 2007 <u>view definition</u>	Crude Rate (CI)	~	<b>1.1</b> (1.1-1.2)
	Age-adjusted Rate (CI)	~	<b>1.1</b> (1.1-1.1)
	Number	319	256,153
Mortality with chronic obstructive pulmonary disease among adults aged 45 years and older - 2007 view definition	Crude Rate (CI)	<b>138.9</b> (123.6-154.1)	<b>223.8</b> (222.9-224.6)
2007 <u>Hew definition</u>	Age-adjusted Rate (CI)	<b>235.2</b> (209.3-261.0)	<b>233.4</b> (232.5-234.3)
Influenza vaccination among adults aged >= 50 years - 2010 view definition	Prevalence (CI)	<b>48.9%</b> (44.5-53.4)	55.0% <sup>a</sup>
Pneumococcal vaccination among adults aged >= 65 years - 2010 view definition	Prevalence (CI)	<b>66.5%</b> (58.8-74.2)	68.8% <sup>a</sup>
	Number	89	90,118
Mortality with end-stage renal disease - 2009 <u>view definition</u>	Crude Rate (CI)	<b>12.7</b> (10.1-15.4)	<b>28.9</b> (28.8-29.1)
	Age-adjusted Rate (CI)	<b>17.7</b> (13.8-21.7)	<b>26.9</b> (26.7-27.0)
	Number	118	114,215
Incidence of treated end-stage renal disease - 2009 <u>view definition</u>	Age-adjusted Rate (CI)	217.9	355.4
	Number	59	49,873
Incidence of treated end-stage renal disease attributed to diabetes - 2009 view definition	Age-adjusted Rate (CI)	104.5	154.1
All teeth lost among adults aged >= 65 years - 2008 view definition	Prevalence (CI)	<b>20.7%</b> (14.9-26.5)	18.5% <sup>a</sup>
Teeth cleaning among adults aged >= 18 years - 2008 <u>view definition</u>	Prevalence	66.3%	71.3% <sup>a</sup>

	(CI)	(63.4-69.2)	
Visits to dentist or dental clinic among adults aged >= 18 years - 2008 view definition	Prevalence (CI)	<b>65.3%</b> (62.4-68.2)	70.0% <sup>a</sup>
Hospitalization for hip fracture among Medicare-eligible persons aged >= 65 years - 2006 view definition	Number	242	234,549
	Crude Rate (CI)	<b>5.6</b> (4.9-6.3)	<b>8.3</b> (8.2-8.3)
Tew demines.	Age-adjusted Rate (CI)	<b>6.5</b> (5.7-7.3)	<b>8.2</b> (8.1-8.2)
	<u> </u>	,	

Note: Numbers presented on this site may differ somewhat from those published by individual states or other federal sources due to differences in methodologies and data sources. The data download does not include footnotes; we recommend printing a print-friendly version of the page to accompany your downloaded data.

## Footnotes

а	Median %.
b	U.S. value includes 11 states.
-	No data available further details may be available from the data source: click on 'view definition' and refer to the link in Data Resources.
~	Data not shown because of too few respondents or cases further details may be available from the data source: click on 'view definition' and refer to the link in Data Resources.
^	Rates are suppressed at state's request.
†	Metropolitan area is defined by multiple counties, but not all counties have data collected.
#	Data were not collected at the county/metropolitan area level.
§	County/Metropolitan area results are not available because the results did not meet program requirements for reporting.
**	Because of the impact from Hurricane's Katrina and Rita on Louisiana's population for the July - December 2005 time period, SEER excluded Louisiana cases diagnosed for that six month time period. Therefore, the incidence count has been suppressed due to data consistency issues. Louisiana incidence rates for 2004-2008 include counts for 2005.

## Measure Definitions

leasure Definitions		
Number	The measured or estimated number of people who had or developed an attribute or disease during a specific year.	
Prevalence	The measured or estimated percentage of people weighted to population characteristics who actually had that attribute or disease during a specific year.	
Age-adjusted Prevalence	Prevalence (see above), standardized to the age distribution of a specific population, usually the standard U.S. 2000 population.	
Crude Rate	The measured number of deaths, cases of disease, or hospitalizations during a specific year – incidence and mortality rates are per 100,000 persons; hospitalization rates are per 1,000 persons	
Age-adjusted Rate	Crude rate (see above) standardized to the age distribution of a specific population, usually the standard U.S. 2000 population.	
Packs Sold per Capita	Packs sold per capita per year	
Mean Days	Weighted to population characteristics	
Years	Years	
CI	95% Confidence Interval.	

http://www.cdc.gov/nccdphp/cdi