



Home > CANSIM Table **577-0013** 1, 2, 3, 4, 5, 6

Aboriginal peoples survey, perceived mental health and suicidal thoughts, by Aboriginal identity, age group and sex, population aged 18 years and over, Canada, provinces and territories

| Data table Add/Remove data ne data below is a part of CANSII | Manipulate Download M table 577-0013. Use the | Related info | rmation Help e data tab to customize your table. | |
|--|--|--|---|--------------------|
| Selected items [Add/Remov | e data] | | | |
| Geography = Yukon Statistics 16, 17, 18 = Number of P | | | | |
| Aboriginal identity 6 | Age group | Sex | Perceived mental health and suicidal thoughts | 2012 |
| Total , Aboriginal identity | Total, 18 years and | | Total, perceived mental health 4, 2 | 3,450 |
| | over | | Excellent or very good perceived mental health 4.2.10 | 1,770 |
| | | | Good perceived mental health 4:2:11 | 1,130 |
| | | | Fair or poor perceived mental health 4: 2: 12 | 400 ^E |
| | | | Perceived mental health not specified ^{4,2,13} | F |
| | | | Total, suicidal thoughts ^{5, 14} | 3,450 |
| | | Both | Ever seriously considered committing suicide 5, 14, 15 | 620 ^E |
| | | sexes | Seriously considered committing suicide in the past 12 months 5, 14, 15 | F |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 15, 15} | 500 ^E |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{2, 13, 14, 15} | F |
| | | | Never seriously considered committing suicide 5. 14 | 2,630 |
| | | Not specified, seriously considered committing suicide 5, 13, 14 | 190 ^l | |
| | | | Total, perceived mental health 4.2 | 1,360 |
| | | | Excellent or very good perceived mental health 4.2.10 | 720 ^l |
| | | | Good perceived mental health 4.2.11 | 490 ^l |
| | | | Fair or poor perceived mental health 4-2-12 | F |
| | | | Perceived mental health not specified 4. 2. 13 | ı |
| | | | Total, suicidal thoughts 5. 14 | 1,360 |
| | | Male | Ever seriously considered committing suicide 5, 14, 15 | 190 |
| | | | Seriously considered committing suicide in the past 12 months 5 14 15 | F |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 14, 15} | 160 ^E |
| | | | Not specified, seriously considered committing suicide in the past 12 months 5 13 14 15 | F |
| | | | Never seriously considered committing suicide 5, 14 | 1,060 |
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | F |
| | | Female | Total, perceived mental health 4.2 | 2,080 |
| | | | Excellent or very good perceived mental health 4.2.10 | 1,050 ^l |
| | | | Good perceived mental health 4: 2:11 | 640 ^l |
| | | | Fair or poor perceived mental health 4, 2, 12 | 300 ^E |
| | | | Perceived mental health not specified 4.2.13 | F |
| | | | Total, suicidal thoughts 5-14 | 2,080 |
| | | | Ever seriously considered committing suicide 5, 14, 15 | 440 |
| | | | Seriously considered committing suicide in the past 12 months 5.15.15 | F |
| | | 1 | | |

| boriginal identity ⁶ | Age group | Sex | Perceived mental health and suicidal thoughts | 201 |
|---------------------------------|-----------------|--------|--|-----|
| | | | Seriously considered committing suicide but not in the past 12 months 5, 14, 15 | 35 |
| | | | Not specified, seriously considered committing suicide in the past 12 months \$\frac{5}{2}\$, \$\frac{13}{2}\$, \$\frac{15}{2}\$ | |
| | | | Never seriously considered committing suicide 5, 14 | 1,5 |
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | |
| | | | Total, perceived mental health 4· 2 | 58 |
| | | | Excellent or very good perceived mental health 4.2.10 | 29 |
| | | | Good perceived mental health 4.9.11 | 20 |
| | | | Fair or poor perceived mental health 4.2.12 | |
| | | | Perceived mental health not specified 4.2.13 | |
| | | | Total, suicidal thoughts ⁵ . ¹⁴ | 5 |
| | | Both | Ever seriously considered committing suicide 5. 15. 15. | |
| | | sexes | Seriously considered committing suicide in the past 12 months 5 14 15 | |
| | | | Seriously considered committing suicide but not in the past 12 months ^{2, 14, 15} | |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 14, 15} | |
| | | | Never seriously considered committing suicide 5 14 | 4 |
| | | | Not specified, seriously considered committing suicide 5. 13. 14 | |
| | | | Total, perceived mental health 4.2 | 2 |
| | | | Excellent or very good perceived mental health 4.2.10 | 1 |
| | | | Good perceived mental health 4.2.11 | |
| | | | Fair or poor perceived mental health 4: 2: 12 | |
| | | | Perceived mental health not specified 4, 2, 13 | |
| | | | Total, suicidal thoughts 5.14 | 2 |
| | 18 to 24 years | Male | Ever seriously considered committing suicide 5, 14, 15 | |
| | 10 10 2 1 yours | Maio | Seriously considered committing suicide in the past 12 months | |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 14, 15} | |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 14, 15} | |
| | | | Never seriously considered committing suicide ⁵ ¹⁴ | 1 |
| | | | Not specified, seriously considered committing suicide ^{5, 13, 14} | |
| | | | Total, perceived mental health 4.2 | 3 |
| | | | Excellent or very good perceived mental health 4.2.19 | |
| | | | Good perceived mental health 4.2.11 | |
| | | | Fair or poor perceived mental health 4. 2. 12 | |
| | | | Perceived mental health not specified 4, 2, 13 | |
| | | | Total, suicidal thoughts 5-14 | 3 |
| | | Female | Ever seriously considered committing suicide 5, 14, 15 | |
| | | | Seriously considered committing suicide in the past 12 months 2.14.15 | |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 14, 15} | |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 14, 15} | |
| | | | Never seriously considered committing suicide \$\frac{5}{2}\$ 14 | 2 |
| | | | Not specified, seriously considered committing suicide ^{5, 13, 14} | |

| ooriginal identity ⁶ | Age group | Sex | Perceived mental health and suicidal thoughts | 2012 |
|---------------------------------|---------------------------|---------------|---|-------|
| | Age group 25 to 54 years | | Total, perceived mental health 4-2 | |
| | | | Excellent or very good perceived mental health 4.2.10 | 1,000 |
| | | | Good perceived mental health 4.2.11 | 610 |
| | | | Fair or poor perceived mental health 4.9.12 | ı |
| | | | Perceived mental health not specified 4.2.13 | |
| | | | Total, suicidal thoughts ^{5, 14} | 1,950 |
| | | Both | Ever seriously considered committing suicide 5, 14, 15 | 380 |
| | | sexes | Seriously considered committing suicide in the past 12 months 5 14 15 | F |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 14, 15} | 300 |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 14, 15} | ı |
| | | | Never seriously considered committing suicide 5-14 | 1,410 |
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | |
| | | | Total, perceived mental health 4.2 | 80 |
| | | | Excellent or very good perceived mental health 4.2, 10 | 450 |
| | | | Good perceived mental health 4.2.11 | 260 |
| | | | Fair or poor perceived mental health 4.2.12 | |
| | | | Perceived mental health not specified 4-2-13 | |
| | | | Total, suicidal thoughts ^{5, 14} | 80 |
| | | | Ever seriously considered committing suicide 5-16-15 | 100 |
| | 25 to 54 years | Male | Seriously considered committing suicide in the past 12 months 5 14 15 | |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 14, 15} | |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 14, 15} | |
| | | | Never seriously considered committing suicide 5-14 | 600 |
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | |
| | | | Total, perceived mental health 4.2 | 1,15 |
| | | | Excellent or very good perceived mental health 4.2.19 | 560 |
| | | | Good perceived mental health 4.2.11 | 350 |
| | | | Fair or poor perceived mental health 4.2.12 | |
| | | | Perceived mental health not specified 4-2-13 | |
| | | | Total, suicidal thoughts ^{5, 14} | 1,15 |
| | | | Ever seriously considered committing suicide 5, 15, 15 | 290 |
| | | Female | Seriously considered committing suicide in the past 12 months | |
| | | | Seriously considered committing suicide but not in the past 12 | 230 |
| | | | months 5 14 15 Not specified, seriously considered committing suicide in the past | 250 |
| | | | 12 months ^{5, 13, 14, 15} | 810 |
| | | | Never seriously considered committing suicide ^{5,14} | |
| | | | Not specified, seriously considered committing suicide ^{5, 13, 14} | 0.10 |
| | 55 years and over | Both sexes | Total, perceived mental health 4.2 | 910 |
| | | | Excellent or very good perceived mental health 4.2.10 | 480 |
| | | | Good perceived mental health 4. 2. 11 | |
| | | | Fair or poor perceived mental health 4.2.12 | |
| | | | Perceived mental health not specified 4, 2, 13 | |

| Aboriginal identity ⁶ | Age group | Sex | Perceived mental health and suicidal thoughts Total, suicidal thoughts 5-14 | 201 |
|----------------------------------|---------------------|--------|---|-----|
| | | | Ever seriously considered committing suicide 5, 14, 15 | |
| | | | Seriously considered committing suicide in the past 12 months 5 14 15 | |
| | | | Seriously considered committing suicide but not in the past 12 months \$\frac{14}{25}\$.15 | |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 14, 15} | |
| | | | Never seriously considered committing suicide 5-14 | 75 |
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | |
| | | | Total, perceived mental health 4.2 | |
| | | | Excellent or very good perceived mental health 4.2.19 | |
| | | | Good perceived mental health 4.2.11 | |
| | | | Fair or poor perceived mental health 4.2.12 | |
| | | | Perceived mental health not specified 4-2-13 | |
| | | | Total, suicidal thoughts ^{5, 14} | |
| | | 24-1- | Ever seriously considered committing suicide 5.14.15 | |
| | | Male | Seriously considered committing suicide in the past 12 months | |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 14, 15} | |
| | | | Not specified, seriously considered committing suicide in the past 12 months 5-12-14-15 | |
| | | | Never seriously considered committing suicide 5-14 | |
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | |
| | | | Total, perceived mental health 4.2 | 6 |
| | | | Excellent or very good perceived mental health 4.2.10 | : |
| | | | Good perceived mental health 4.2.11 | |
| | | | Fair or poor perceived mental health 4-2-12 | |
| | | | Perceived mental health not specified 4-2-13 | |
| | | | Total, suicidal thoughts ^{5, 14} | |
| | | Female | Ever seriously considered committing suicide 5.16.15 | |
| | | Female | Seriously considered committing suicide in the past 12 months 5, 14, 15 | |
| | | | Seriously considered committing suicide but not in the past 12 months 5, 16, 15 | |
| | | | Not specified, seriously considered committing suicide in the past 12 months \$1.15.15 | |
| | | | Never seriously considered committing suicide 5, 14 | 4 |
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | |
| st Nations (North American | Total, 18 years and | Both | Total, perceived mental health 4.2 | 2, |
| dian) | over | sexes | Excellent or very good perceived mental health 4.2.10 | 1, |
| | | | Good perceived mental health 4: 2: 11 | 8 |
| | | | Fair or poor perceived mental health 4.2.12 | : |
| | | | Perceived mental health not specified 4.2.13 | |
| | | | Total, suicidal thoughts ^{5, 14} | 2, |
| | | | Ever seriously considered committing suicide 5, 14, 15 | |
| | | | Seriously considered committing suicide in the past 12 months 5, 14, 15 | |
| | | | Seriously considered committing suicide but not in the past 12 months \$\frac{14.15}{2}\$ | 3 |

| Aboriginal identity ⁶ | Age group | Sex | Perceived mental health and suicidal thoughts | 2012 |
|----------------------------------|---------------------------|---------------|---|-------|
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 16} | |
| | Age group 18 to 24 years | | Never seriously considered committing suicide ^{5, 14} | 2,15 |
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | |
| | | | Total, perceived mental health 4.2 | 1,120 |
| | | | Excellent or very good perceived mental health 4.2.10 | 610 |
| | | | Good perceived mental health 4. 2. 11. | 410 |
| | | | Fair or poor perceived mental health 4.2.12 | |
| | | | Perceived mental health not specified 4-2-13 | |
| | | | Total, suicidal thoughts 5 14 | 1,12 |
| | | Male | Ever seriously considered committing suicide 5, 14, 15 | 16 |
| | | | Seriously considered committing suicide in the past 12 months 5.14.15 | |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 14, 15} | |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 16, 15} | |
| | | | Never seriously considered committing suicide ^{5, 14} | 89 |
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | |
| | | | Total, perceived mental health 4, 2 | 1,6 |
| | | | Excellent or very good perceived mental health 4.2.10 | 92 |
| | | | Good perceived mental health 4. 2.11 | 44 |
| | | | Fair or poor perceived mental health 4.2.12 | |
| | | | Perceived mental health not specified 4-2-13 | |
| | | | Total, suicidal thoughts 5.14 | 1,6 |
| | | Female | Ever seriously considered committing suicide 5, 14, 15 | 35 |
| | | | Seriously considered committing suicide in the past 12 months 5-14-15 | |
| | | | Seriously considered committing suicide but not in the past 12 months 5-14-15 | 26 |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{2, 12, 16, 18} | |
| | | | Never seriously considered committing suicide 5.14 | 1,2 |
| | | | Not specified, seriously considered committing suicide 5-13-14 | |
| | 18 to 24 years | | Total, perceived mental health 4.2 | 49 |
| | | | Excellent or very good perceived mental health 4.2.10 | 26 |
| | | | Good perceived mental health 4, 2, 11 | 15 |
| | | | Fair or poor perceived mental health 4.2.12 | |
| | | | Perceived mental health not specified 4-2-13 | |
| | | | Total, suicidal thoughts ^{5, 14} | 49 |
| | | Both sexes | Ever seriously considered committing suicide 5, 14, 15 | |
| | | Sexes | Seriously considered committing suicide in the past 12 months 5-14-15 | |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 14, 15} | |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 16, 15} | |
| | | | Never seriously considered committing suicide ⁵ · ¹⁴ | 39 |
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | |
| | | Male | Total, perceived mental health 4-2 | 22 |
| | | | Excellent or very good perceived mental health 4. 2, 10 | 10 |

| boriginal identity ⁶ | Age group | Sex | Perceived mental health and suicidal thoughts | 2012 |
|---------------------------------|----------------|--------|--|------|
| | | | Good perceived mental health 4.2.11 | |
| | | | Fair or poor perceived mental health 4.2.12 | |
| | | | Perceived mental health not specified 4.2.13 | |
| | | | Total, suicidal thoughts 5-14 | 220 |
| | | | Ever seriously considered committing suicide 5- 14- 15 | |
| | | | Seriously considered committing suicide in the past 12 months | |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 12, 15} | |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{2, 12, 16, 15} | |
| | | | Never seriously considered committing suicide 5.14 | 150 |
| | | | Not specified, seriously considered committing suicide 5. 13. 14 | |
| | | | Total, perceived mental health 4.2 | 27 |
| | | | Excellent or very good perceived mental health 4: 2: 10 | |
| | | | Good perceived mental health 4.2.11 | |
| | | | Fair or poor perceived mental health 4.2.12 | |
| | | | Perceived mental health not specified 4, 2, 13 | |
| | | | Total, suicidal thoughts 5, 14 | 27 |
| | | Female | Ever seriously considered committing suicide 5, 14, 15 | |
| | | | Seriously considered committing suicide in the past 12 months 5-14-15 | |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 14, 15} | |
| | | | Not specified, seriously considered committing suicide in the past 12 months $^{5\cdot13\cdot14\cdot15}$ | |
| | | | Never seriously considered committing suicide 5.14 | 24 |
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | |
| | 25 to 54 years | | Total, perceived mental health 4.2 | 1,5 |
| | | | Excellent or very good perceived mental health 4.2.10 | 87 |
| | | | Good perceived mental health 4.2.11 | 42 |
| | | | Fair or poor perceived mental health 4-2-12 | |
| | | | Perceived mental health not specified 4, 2, 13 | |
| | | | Total, suicidal thoughts 5.14 | 1,5 |
| | | Both | Ever seriously considered committing suicide 5, 14, 15 | 29 |
| | | sexes | Seriously considered committing suicide in the past 12 months 2 14 15 | |
| | | | Seriously considered committing suicide but not in the past 12 months ^{3, 14, 15} | 20 |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 14, 15} | |
| | | | Never seriously considered committing suicide ^{5, 14} | 1,1 |
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | |
| | | Male | Total, perceived mental health 4. 2 | 66 |
| | | | Excellent or very good perceived mental health 4 2 10 | 39 |
| | | | Good perceived mental health 4.2.11 | 22 |
| | | | Fair or poor perceived mental health 4.2.12 | |
| | | | Perceived mental health not specified 4.2.13 | |
| | | | Total, suicidal thoughts ^{5, 14} | 66 |
| | | | Ever seriously considered committing suicide \$\frac{5}{2}\$. 16. 15 | |

| Aboriginal identity 6 | Age group | Sex | Perceived mental health and suicidal thoughts | 2012 |
|-----------------------|-------------------|--------|--|------|
| | | | Seriously considered committing suicide in the past 12 months 5 14 15 | |
| | | | Seriously considered committing suicide but not in the past 12 months ⁵ ¹⁵ ¹⁵ | 1 |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 14, 15} | ı |
| | | | Never seriously considered committing suicide ^{5, 14} | 500 |
| | | | Not specified, seriously considered committing suicide ^{5, 13, 14} | |
| | | | Total, perceived mental health 4.2 | 880 |
| | | | Excellent or very good perceived mental health 4.2.19 | 480 |
| | | | Good perceived mental health 4.2.11 | |
| | | | Fair or poor perceived mental health 4: 2: 12 | |
| | | | Perceived mental health not specified 4, 2, 13 | |
| | | | Total, suicidal thoughts 5. 14 | 880 |
| | | Female | Ever seriously considered committing suicide 5, 14, 15 | 210 |
| | | | Seriously considered committing suicide in the past 12 months 5 14 15 | |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 15, 15} | |
| | | | Not specified, seriously considered committing suicide in the past 12 months 5 13 14 15 | |
| | | | Never seriously considered committing suicide ^{5, 14} | 620 |
| | | | Not specified, seriously considered committing suicide ^{5, 13, 14} | |
| | 55 years and over | | Total, perceived mental health 4-2 | 77 |
| | | | Excellent or very good perceived mental health 4.2.10 | 40 |
| | | | Good perceived mental health 4· 2· 11 | |
| | | | Fair or poor perceived mental health 4-2-12 | |
| | | | Perceived mental health not specified 4, 2, 13 | |
| | | | Total, suicidal thoughts 5. 14 | 77 |
| | | Both | Ever seriously considered committing suicide 5, 14, 15 | |
| | | sexes | Seriously considered committing suicide in the past 12 months 5.14.15 | |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 16, 15} | |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 14, 15} | |
| | | | Never seriously considered committing suicide 5 14 | 65 |
| | | | Not specified, seriously considered committing suicide ^{5, 13, 14} | |
| | | Male | Total, perceived mental health 4.2 | |
| | | | Excellent or very good perceived mental health 4.2.10 | |
| | | | Good perceived mental health 4.2.11 | |
| | | | Fair or poor perceived mental health 4, 2, 12 | |
| | | | Perceived mental health not specified 4, 2, 13 | |
| | | | Total, suicidal thoughts ^{5, 14} | |
| | | | Ever seriously considered committing suicide 5, 14, 15 | |
| | | | Seriously considered committing suicide in the past 12 months 5.14.15 | |
| | | | Seriously considered committing suicide but not in the past 12 months ^{2, 14, 15} | |
| | | | Not specified, seriously considered committing suicide in the past 12 months 5, 13, 14, 15 | |
| | | | Never seriously considered committing suicide 5, 14 | |

| Aboriginal identity 6 | Age group | Sex | Perceived mental health and suicidal thoughts | 2012 |
|-----------------------|--------------------------|--------|---|------------------|
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | F |
| | | | Total, perceived mental health 4.2 | 530 ^E |
| | | | Excellent or very good perceived mental health 4-2-10 | F |
| | | | Good perceived mental health 4.2.11 | F |
| | | | Fair or poor perceived mental health 4, 2, 12 | F |
| | | | Perceived mental health not specified 4, 2, 13 | F |
| | | | Total, suicidal thoughts ^{5, 14} | 530 ^E |
| | | Female | Ever seriously considered committing suicide 5, 14, 15 | F |
| | | | Seriously considered committing suicide in the past 12 months 5 14 15 | F |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 14, 15} | F |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 14, 15} | F |
| | | | Never seriously considered committing suicide 5, 14 | 420 ^E |
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | F |
| Métis | Total, 18 years and over | | Total, perceived mental health 4: 2 | 580 |
| | 0001 | | Excellent or very good perceived mental health 4.2.10 | 220 ^E |
| | | | Good perceived mental health 4.2.11 | 240 ^E |
| | | | Fair or poor perceived mental health 4.2.12 | F |
| | | | Perceived mental health not specified 4: 2: 13 | F |
| | | | Total, suicidal thoughts ^{5, 14} | 580 |
| | | Both | Ever seriously considered committing suicide 5, 14, 15 | F |
| | | sexes | Seriously considered committing suicide in the past 12 months 5.16.15 | F |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 14, 15} | F |
| | | | Not specified, seriously considered committing suicide in the past 12 months ²⁻¹³⁻¹⁵ | F |
| | | | Never seriously considered committing suicide 5, 14 | 430 |
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | F |
| | | | Total, perceived mental health 4.2 | 230 ^E |
| | | | Excellent or very good perceived mental health 4.2.19 | 110 ^E |
| | | | Good perceived mental health 4.2.11 | F |
| | | | Fair or poor perceived mental health 4. 2. 12 | F |
| | | | Perceived mental health not specified 4. 2. 13 | F |
| | | | Total, suicidal thoughts ^{5, 14} | 230 ^E |
| | | Male | Ever seriously considered committing suicide ^{5, 14, 15} | F |
| | | | Seriously considered committing suicide in the past 12 months 5.14.15 | F |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 16, 15} | F |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 14, 15} | F |
| | | | Never seriously considered committing suicide 5-14 | 160 ^E |
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | F |
| | | Female | Total, perceived mental health 4· 2 | 350 ^E |
| | | | Excellent or very good perceived mental health 4.2.10 | 110 ^E |
| | | | Good perceived mental health 4.2.11 | 180 ^E |
| | | | Fair or poor perceived mental health 4: 2: 12 | F |

| boriginal identity ⁶ | Age group | Sex | Perceived mental health and suicidal thoughts | 2012 |
|---------------------------------|----------------|--------|--|------|
| | | | Perceived mental health not specified 4.2.13 | |
| | | | Total, suicidal thoughts ^{5, 14} | 350 |
| | | | Ever seriously considered committing suicide ^{5, 16, 15} | |
| | | | Seriously considered committing suicide in the past 12 months 5.14.15 | |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 15, 15} | |
| | | | Not specified, seriously considered committing suicide in the past 12 months $^{\frac{5}{12},\frac{13}{15},\frac{15}{15}}$ | |
| | | | Never seriously considered committing suicide 5.14 | 270 |
| | | | Not specified, seriously considered committing suicide ^{5, 13, 14} | |
| | 18 to 24 years | | Total, perceived mental health 4.2 | |
| | | | Excellent or very good perceived mental health 4.2.10 | |
| | | | Good perceived mental health 4.2.11 | |
| | | | Fair or poor perceived mental health 4: 9: 12 | |
| | | | Perceived mental health not specified 4, 2, 13 | |
| | | | Total, suicidal thoughts ^{5, 14} | |
| | | Both | Ever seriously considered committing suicide 5, 14, 15 | |
| | | sexes | Seriously considered committing suicide in the past 12 months 5.14.15 | |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 14, 15} | |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 12, 14, 15} | |
| | | | Never seriously considered committing suicide 5 14 | |
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | |
| | | | Total, perceived mental health 4.2 | |
| | | | Excellent or very good perceived mental health 4.2.19 | |
| | | | Good perceived mental health 4.2.11 | |
| | | | Fair or poor perceived mental health 4.2.12 | |
| | | | Perceived mental health not specified 4, 2, 13 | |
| | | | Total, suicidal thoughts 5. 14 | |
| | | Male | Ever seriously considered committing suicide 5, 14, 15 | |
| | | | Seriously considered committing suicide in the past 12 months 5.14.15 | |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 14, 15} | |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 14, 15} | |
| | | | Never seriously considered committing suicide ^{5, 14} | |
| | | | Not specified, seriously considered committing suicide ^{5, 13, 14} | |
| | | Female | Total, perceived mental health 4.2 | |
| | | | Excellent or very good perceived mental health 4.9.10 | |
| | | | Good perceived mental health 4.2.11 | |
| | | | Fair or poor perceived mental health 4.2.12 | |
| | | | Perceived mental health not specified 4. 2. 13 | |
| | | | Total, suicidal thoughts ^{5, 14} | |
| | | | Ever seriously considered committing suicide 5, 14, 15 | |
| | | | Seriously considered committing suicide in the past 12 months 5 15 15 15 | |
| | | | | |

| boriginal identity ⁶ | Age group | Sex | Perceived mental health and suicidal thoughts | 201 |
|---------------------------------|----------------|--------|--|-----|
| | | | Seriously considered committing suicide but not in the past 12 months 5, 14, 15 | |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{2, 13, 14, 15} | |
| | | | Never seriously considered committing suicide 5, 14 | |
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | |
| | | | Total, perceived mental health 4· 2 | 400 |
| | | | Excellent or very good perceived mental health 4.2.10 | 130 |
| | | | Good perceived mental health 4.2.11 | 18 |
| | | | Fair or poor perceived mental health 4.2.12 | |
| | | | Perceived mental health not specified 4.2.13 | |
| | | | Total, suicidal thoughts ^{5, 14} | 40 |
| | | Both | Ever seriously considered committing suicide 5.14.15 | |
| | | sexes | Seriously considered committing suicide in the past 12 months 5-14-15 | |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 14, 15} | |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 14, 15} | |
| | | | Never seriously considered committing suicide 5-14 | 28 |
| | | | Not specified, seriously considered committing suicide ^{5, 12, 14} | |
| | | | Total, perceived mental health 4, 2 | 1: |
| | | | Excellent or very good perceived mental health 4.2.10 | |
| | | | Good perceived mental health 4.2.11 | |
| | | | Fair or poor perceived mental health 4.2.12 | |
| | | | Perceived mental health not specified 4.2.13 | |
| | | | Total, suicidal thoughts 5-14 | 1 |
| | 25 to 54 years | Male | Ever seriously considered committing suicide 5, 14, 15 | |
| | | | Seriously considered committing suicide in the past 12 months 5.16.15 | |
| | | | Seriously considered committing suicide but not in the past 12 months 5-14-15 | |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 14, 15} | |
| | | | Never seriously considered committing suicide ^{5, 14} | |
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | |
| | | | Total, perceived mental health 4, 2 | 2 |
| | | | Excellent or very good perceived mental health 4.9.10 | |
| | | | Good perceived mental health 4.2.11 | |
| | | | Fair or poor perceived mental health 4·2·12 | |
| | | | Perceived mental health not specified 4, 2, 13 | |
| | | | Total, suicidal thoughts ^{5, 14} | 2 |
| | | Female | Ever seriously considered committing suicide 5, 14, 15 | |
| | | | Seriously considered committing suicide in the past 12 months 5-15-15 | |
| | | | Seriously considered committing suicide but not in the past 12 months ⁵ ¹⁴ ¹⁵ | |
| | | | Not specified, seriously considered committing suicide in the past 12 months 5 13 16 15 | |
| | | | Never seriously considered committing suicide 5-14 | 19 |
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | |

| boriginal identity ⁶ | Age group | Sex | Perceived mental health and suicidal thoughts | 20 |
|---------------------------------|--------------------------|---------------|--|----|
| | | | Total, perceived mental health 4.2 | |
| | | | Excellent or very good perceived mental health 4-2-10 | |
| | | | Good perceived mental health 4, 2, 11 | |
| | | | Fair or poor perceived mental health 4:2:12 | |
| | | | Perceived mental health not specified 4.2.13 | |
| | | | Total, suicidal thoughts 5-14 | |
| | | Both | Ever seriously considered committing suicide 5, 14, 15 | |
| | | sexes | Seriously considered committing suicide in the past 12 months 5.14.15 | |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 16, 15} | |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 14, 15} | |
| | | | Never seriously considered committing suicide 5, 14 | |
| | | | Not specified, seriously considered committing suicide ^{5, 13, 14} | |
| | | | Total, perceived mental health 4.2 | |
| | | | Excellent or very good perceived mental health 4.2.10 | |
| | | | Good perceived mental health 4.2.11 | + |
| | | | Fair or poor perceived mental health 4.2.12 | + |
| | | | Perceived mental health not specified 4.2.13 | + |
| | | | Total, suicidal thoughts ^{5, 14} | + |
| | | | Ever seriously considered committing suicide \$\frac{5}{2}\$. 15 | + |
| | 55 years and over | Male | Seriously considered committing suicide in the past 12 months | |
| | | | 5, 14, 15 | |
| | | | Seriously considered committing suicide but not in the past 12 months 2 12 15 | |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 12, 14, 15} | |
| | | | Never seriously considered committing suicide 5.14 | |
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | |
| | | | Total, perceived mental health 4.2 | |
| | | | Excellent or very good perceived mental health 4.2.10 | |
| | | | Good perceived mental health 4.2.11 | |
| | | | Fair or poor perceived mental health 4-2-12 | |
| | | | Perceived mental health not specified 4-2-13 | |
| | | | Total, suicidal thoughts ^{5, 14} | + |
| | | | Ever seriously considered committing suicide 5, 14, 15 | + |
| | | Female | Seriously considered committing suicide in the past 12 months | + |
| | | | Seriously considered committing suicide but not in the past 12 | |
| | | | months ⁵ . ^{14. 15} Not specified, seriously considered committing suicide in the past | |
| | | | 12 months ^{5, 13, 14, 15} | - |
| | | | Never seriously considered committing suicide 5-14 | - |
| | | | Not specified, seriously considered committing suicide 5: 13, 14 | _ |
| uk (Inuit) | Total, 18 years and over | Both sexes | Total, perceived mental health 4.2 | |
| | | | Excellent or very good perceived mental health 4 2 10 | |
| | | | Good perceived mental health 4.9.11 | |
| | | | Fair or poor perceived mental health 4.9.12 | |
| | | | Perceived mental health not specified 4.2.13 | |

| iginal identity ⁶ | Age group | Sex | Perceived mental health and suicidal thoughts | 2012 |
|------------------------------|----------------|--------|---|------|
| | | | Total, suicidal thoughts ^{5, 14} | F |
| | | | Ever seriously considered committing suicide 5. 15. 15. | F |
| | | | Seriously considered committing suicide in the past 12 months | |
| | | | Seriously considered committing suicide but not in the past 12 months ⁵⁻¹⁴⁻¹⁵ | F |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 16, 18} | F |
| | | | Never seriously considered committing suicide 5.14 | F |
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | F |
| | | | Total, perceived mental health 4.2 | F |
| | | | Excellent or very good perceived mental health 4.2.10 | F |
| | | | Good perceived mental health 4. 2.11 | F |
| | | | Fair or poor perceived mental health 4.2.12 | F |
| | | | Perceived mental health not specified 4, 2, 13 | F |
| | | | Total, suicidal thoughts 5, 14 | F |
| | | Male | Ever seriously considered committing suicide ^{5, 14, 15} | F |
| | | | Seriously considered committing suicide in the past 12 months 5-15-15 | F |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 14, 15} | F |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 14, 15} | F |
| | | | Never seriously considered committing suicide 5.14 | F |
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | F |
| | | | Total, perceived mental health 4.2 | F |
| | | | Excellent or very good perceived mental health 4.2.10 | F |
| | | | Good perceived mental health 4. 2. 11 | F |
| | | | Fair or poor perceived mental health 4.2.12 | F |
| | | | Perceived mental health not specified 4.2.13 | F |
| | | | Total, suicidal thoughts 5 14 | F |
| | | Female | Ever seriously considered committing suicide 5. 16. 15 | F |
| | | | Seriously considered committing suicide in the past 12 months 5 14 15 | F |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 14, 15} | F |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{2, 13, 16, 15} | F |
| | | | Never seriously considered committing suicide 5.14 | F |
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | F |
| | 18 to 24 years | Both | Total, perceived mental health 4.2 | F |
| | | sexes | Excellent or very good perceived mental health 4.2.10 | F |
| | | | Good perceived mental health 4, 2, 11 | F |
| | | | Fair or poor perceived mental health 4.2.12 | F |
| | | | Perceived mental health not specified 4-2-13 | F |
| | | | Total, suicidal thoughts ^{5, 14} | F |
| | | | Ever seriously considered committing suicide 5, 14, 15 | F |
| | | | Seriously considered committing suicide in the past 12 months | F |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 14, 15} | F |

| boriginal identity ⁶ | Age group | Sex | Perceived mental health and suicidal thoughts | 201 |
|---------------------------------|----------------|---------------|--|-----|
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 14, 15} | |
| | | | Never seriously considered committing suicide ^{5, 14} | |
| | | | Not specified, seriously considered committing suicide ^{5, 13, 14} | |
| | | Male | Total, perceived mental health 4-2 | |
| | | | Excellent or very good perceived mental health 4.2.10 | |
| | | | Good perceived mental health 4.2.11 | |
| | | | Fair or poor perceived mental health 4: 2: 12 | |
| | | | Perceived mental health not specified 4, 2, 13 | |
| | | | Total, suicidal thoughts 5. 14 | |
| | | | Ever seriously considered committing suicide 5, 14, 15 | |
| | | | Seriously considered committing suicide in the past 12 months 5.14.15 | |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 16, 15} | |
| | | | Not specified, seriously considered committing suicide in the past 12 months $^{5\cdot13\cdot14\cdot15}$ | |
| | | | Never seriously considered committing suicide ^{5, 14} | |
| | | | Not specified, seriously considered committing suicide ^{5, 13, 14} | |
| | | Female | Total, perceived mental health 4.2 | |
| | | | Excellent or very good perceived mental health 4.2.19 | |
| | | | Good perceived mental health 4.2.11 | |
| | | | Fair or poor perceived mental health 4, 9, 12 | |
| | | | Perceived mental health not specified 4, 2, 13 | |
| | | | Total, suicidal thoughts ^{5, 14} | |
| | | | Ever seriously considered committing suicide 5, 14, 15 | |
| | | | Seriously considered committing suicide in the past 12 months 5.14.15 | |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 14, 15} | |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 14, 15} | |
| | | | Never seriously considered committing suicide 5. 14 | |
| | | | Not specified, seriously considered committing suicide ^{5, 13, 14} | |
| | 25 to 54 years | Both sexes | Total, perceived mental health 4, 2 | |
| | | | Excellent or very good perceived mental health 4.2.19 | |
| | | | Good perceived mental health 4.2.11 | |
| | | | Fair or poor perceived mental health 4. 2. 12 | |
| | | | Perceived mental health not specified 4, 2, 13 | |
| | | | Total, suicidal thoughts 5, 14 | |
| | | | Ever seriously considered committing suicide 5, 14, 15 | |
| | | | Seriously considered committing suicide in the past 12 months 5.14.15 | |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 14, 15} | |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 16, 15} | |
| | | | Never seriously considered committing suicide ^{5, 14} | |
| | | | Not specified, seriously considered committing suicide ^{5, 13, 14} | |
| | | Male | Total, perceived mental health 4: 2 | |
| | | | Excellent or very good perceived mental health 4.2.19 | |

| Good perceived mental health * 2 ± 12 Fair or poor perceived mental health * 2 ± 14 Perceived mental health not specified * 2 ± 14 Total, suicidal thoughts * 12 Ever seriously considered committing suicide in the past 12 more * 2 ± 14 Seriously considered committing suicide but not in the past northers * 2 ± 14 Not specified, seriously considered committing suicide in the past northers * 2 ± 14 Not specified, seriously considered committing suicide in the past northers * 2 ± 14 Not specified, seriously considered committing suicide * 2 ± 14 Not specified, seriously considered committing suicide * 2 ± 14 Not specified, seriously considered committing suicide * 2 ± 14 Excellent or very good perceived mental health * 2 ± 14 Excellent or very good perceived mental health * 2 ± 14 Fair or poor perceived mental health * 2 ± 14 Fair or poor perceived mental health * 2 ± 14 Perceived mental health not specified * 2 ± 14 Total, suicidal thoughts * ± 14 Ever seriously considered committing suicide in the past not specified, seriously considered committing suicide in the past not specified, seriously considered committing suicide in the past not specified, seriously considered committing suicide in the past not specified, seriously considered committing suicide in the past not specified, seriously considered committing suicide in the past not specified seriously considered committing suicide in the past not specified seriously considered committing suicide in the past not specified seriously considered committing suicide in the past not specified seriously considered committing suicide in the past not specified seriously considered committing suicide in the past not specified seriously considered committing suicide in the past not specified seriously considered committing suicide in the past not specified seriously considered committing suicide in the past not specified seriously considered committing suicide in the past not specified seriously considered committing suicide in the past not specified seriously considered c | 20 |
|---|------|
| Perceived mental health not specified \$\pi_{\pi_{\pi_{\pi_{\pi_{\pi_{\pi_{\pi_ | |
| Total, suicidal thoughts \$\frac{\pmu}{\pmu}\$ Ever seriously considered committing suicide \$\frac{\pmu}{\pmu}\$ \frac{\pmu}{\pmu}\$ Seriously considered committing suicide in the past 12 mon shall Seriously considered committing suicide but not in the past not \$\pmu\$ seriously considered committing suicide in the \$\pmu\$ seriously considered committing suicide in the \$\pmu\$ seriously considered committing suicide \$\frac{\pmu}{\pmu}\$ Not specified, seriously considered committing suicide \$\frac{\pmu}{\pmu}\$ Not specified, seriously considered committing suicide \$\frac{\pmu}{\pmu}\$ Fair or poor perceived mental health \$\frac{\pmu}{\pmu}\$ Female Female Female Female Female Female Female Seriously considered committing suicide \$\frac{\pmu}{\pmu}\$ Not specified, seriously considered committing suicide \$\frac{\pmu}{\pmu}\$ Not specified, seriously considered committing suicide \$\frac{\pmu}{\pmu}\$ Not specified, seriously considered committing suicide in the past 12 mon \$\frac{\pmu}{\pmu}\$ \frac{\pmu}{\pmu}\$ \pmu | |
| Ever seriously considered committing suicide \$\frac{\f | |
| Seriously considered committing suicide but not in the past months 2 to 2 to 3 to 3 to 3 to 3 to 3 to 3 to | |
| Seriously considered committing suicide but not in the past months \$\frac{2}{12}\$. Not specified, seriously considered committing suicide \$\frac{1}{12}\$. Food of perceived mental health \$\frac{1}{12}\$. Excellent or very good perceived mental health \$\frac{1}{12}\$. Fair or poor perceived mental health \$\frac{1}{12}\$. Fermalo Femalo Femalo Femalo Femalo Femalo Seriously considered committing suicide \$\frac{1}{12}\$. Ever seriously considered committing suicide \$\frac{1}{12}\$. Not specified, seriously considered committing suicide in the past 12 months \$\frac{1}{12}\$. Not specified, seriously considered committing suicide \$\frac{1}{12}\$. Not specified, seriously considered committing suicide \$\frac{1}{12}\$. Not specified, seriously considered committing suicide \$\frac{1}{12}\$. Not specified mental health \$\frac{1}{12}\$. Excellent or very good perceived mental health \$\frac{1}{12}\$. Fair or poor perceived mental health \$\frac{1}{12}\$. Ferceived mental health \$\frac{1}{12}\$. Ferceived mental health \$\frac{1}{12}\$. Ever seriously considered committing suicide \$\frac{1}{12}\$. Fair or poor perceived mental health \$\frac{1}{12}\$. Ferceived mental health \$\frac{1}{12}\$. Seriously considered committing suicide \$\frac{1}{12}\$. Not specified, seriously considered committing suicide in the past 12 months \$\frac{1}{12}\$. Not specified, seriously considered committing suicide in the past 12 months \$\frac{1}{12}\$. Not specified, seriously considered committing suicide in the past nonths \$\frac{1}{12}\$. Not specified, seriously considered committing suicide in the past 12 months \$\frac{1}{12}\$. Not specified, seriously considered committing suicide in the past 12 months \$\frac{1}{12}\$. | ıs |
| months \$\frac{24}{24}\$ Not specified, seriously considered committing suicide in the 2 months \$\frac{1}{24}\$ is \$\frac{1}{24}\$. Never seriously considered committing suicide \$\frac{1}{24}\$ is \$\frac{1}{24}\$. Not specified, seriously considered committing suicide \$\frac{1}{24}\$ is \$\frac{1}{24}\$. Total, perceived mental health \$\frac{1}{2}\$ is \$\frac{1}{24}\$. Excellent or very good perceived mental health \$\frac{1}{2}\$ is \$\frac{1}{24}\$. Fair or poor perceived mental health \$\frac{1}{2}\$ is \$\frac{1}{24}\$. Perceived mental health not specified \$\frac{1}{2}\$ is \$\frac{1}{2}\$. Female Female Female Female Female Female Female Female Seriously considered committing suicide \$\frac{1}{2}\$ is \$\frac{1}{ | |
| Never seriously considered committing suicide \$\partial \partial \par | 2 |
| Not specified, seriously considered committing suicide \$\frac{2}{2} interpretable of the past o | past |
| Total, perceived mental health \$\frac{1}{2}\$ Excellent or very good perceived mental health \$\frac{1}{2}\$ Good perceived mental health \$\frac{1}{2}\$ Fair or poor perceived mental health \$\frac{1}{2}\$ Perceived mental health not specified \$\frac{1}{2}\$ \$\frac{1}{2}\$ Total, suicidal thoughts \$\frac{1}{2}\$ Female | |
| Excellent or very good perceived mental health \$\frac{b}{2} \frac{12}{12}\$ Good perceived mental health \$\frac{b}{2} \frac{12}{12}\$ Fair or poor perceived mental health \$\frac{b}{2} \frac{12}{12}\$ Perceived mental health not specified \$\frac{d}{d} \frac{2}{2} \frac{12}{12}\$ Total, suicidal thoughts \$\frac{b}{2} \frac{12}{12}\$ Female Ever seriously considered committing suicide \$\frac{b}{1} \frac{12}{12}\$ Seriously considered committing suicide \$\frac{b}{1} \frac{12}{12}\$ Seriously considered committing suicide but not in the past nonths \$\frac{b}{2} \frac{b}{1} \frac{b}{2} \f | |
| Female Fever seriously considered committing suicide in the past 12 mon in the past months **± ± ± ± ± ± ± ± ± ± ± ± ± ± ± ± ± ± | |
| Fair or poor perceived mental health \$\frac{2}{2} \frac{12}{2}\$ Perceived mental health not specified \$\frac{2}{2} \frac{12}{2}\$ Total, suicidal thoughts \$\frac{2}{2} \frac{12}{2}\$ Female Ever seriously considered committing suicide \$\frac{2}{2} \frac{12}{2} \frac{12}{2}\$ Seriously considered committing suicide but not in the past nonths \$\frac{2}{2} \frac{12}{2} \frac{12}{2}\$ Not specified, seriously considered committing suicide \$\frac{2}{2} \frac{12}{2}\$ Not specified, seriously considered committing suicide \$\frac{2}{2} \frac{12}{2}\$ Not specified, seriously considered committing suicide \$\frac{2}{2} \frac{12}{2}\$ Excellent or very good perceived mental health \$\frac{2}{2} \frac{12}{2}\$ Fair or poor perceived mental health \$\frac{2}{2} \frac{12}{2}\$ Seriously considered committing suicide \$\frac{2}{2} \frac{12}{2}\$ Seriously considered committing suicide \$\frac{2}{2} \frac{12}{2}\$ Not specified, seriously considered committing suicide \$\frac{2}{2} \frac{12}{2}\$ Perceived mental health \$\frac{2}{2} \frac{12}{2}\$ Not specified, seriously considered committing suicide \$\frac{2}{2} \frac{12}{2}\$ Perceived mental health \$\frac{2}{2} \frac{12}{2}\$ Perceived men | |
| Fair or poor perceived mental health \$\frac{2}{2} \frac{12}{2}\$ Perceived mental health not specified \$\frac{2}{2} \frac{12}{2}\$ Total, suicidal thoughts \$\frac{2}{2} \frac{1}{2}\$ Female Ever seriously considered committing suicide \$\frac{2}{2} \frac{12}{2} \frac{12}{2}\$ Seriously considered committing suicide but not in the past nonths \$\frac{2}{2} \frac{12}{2} \frac{12}{2}\$ Not specified, seriously considered committing suicide in the 12 months \$\frac{2}{2} \frac{12}{2} \frac{12}{2}\$ Not specified, seriously considered committing suicide \$\frac{2}{2} \frac{12}{2}\$ Not specified, seriously considered committing suicide \$\frac{2}{2} \frac{12}{2}\$ Excellent or very good perceived mental health \$\frac{2}{2} \frac{12}{2}\$ Fair or poor perceived mental health \$\frac{2}{2} \frac{12}{2}\$ Seriously considered committing suicide \$\frac{2}{2} \frac{12}{2} \frac{12}{2}\$ Seriously considered committing suicide but not in the past nonths \$\frac{2}{2} \frac{12}{2} \frac{12}{2}\$ Not specified, seriously considered committing suicide in the 12 months \$\frac{2}{2} \frac{12}{2} | |
| Perceived mental health not specified \$\frac{2.21}{2.24}\$ Total, suicidal thoughts \$\frac{2.24}{2.24}\$ Ever seriously considered committing suicide in the past 12 mon \$\frac{2.24}{2.24}\$ Seriously considered committing suicide but not in the past months \$\frac{2.24}{2.24}\$ Not specified, seriously considered committing suicide in the 12 months \$\frac{2.24}{2.24}\$ Not specified, seriously considered committing suicide in the 12 months \$\frac{2.24}{2.24}\$ Not specified, seriously considered committing suicide \$\frac{2.24}{2.24}\$ Not specified, seriously considered committing suicide \$\frac{2.24}{2.24}\$ Not specified, seriously considered committing suicide \$\frac{2.24}{2.24}\$ Excellent or very good perceived mental health \$\frac{4.2}{2.24}\$ Fair or poor perceived mental health \$\frac{4.2}{2.24}\$ Perceived mental health not specified \$\frac{4.2}{2.24}\$ Total, suicidal thoughts \$\frac{2.24}{2.24}\$ Perceived mental health not specified \$\frac{4.2}{2.24}\$ Total, suicidal thoughts \$\frac{2.24}{2.24}\$ Ever seriously considered committing suicide \$\frac{2.24}{2.24}\$ Not specified, seriously considered committing suicide in the 12 months \$\frac{2.24}{2.24}\$ Not specified, seriously considered committing suicide in the 12 months \$\frac{2.24}{2.24}\$ Not specified, seriously considered committing suicide in the 12 months \$\frac{2.24}{2.24}\$ Not specified, seriously considered committing suicide in the 12 months \$\frac{2.24}{2.24}\$ Not specified, seriously considered committing suicide \$\frac{2.24}{2.24}\$ Not specified, seriously considered committing suici | |
| Female Female Ever seriously considered committing suicide \$\frac{1}{2}\fra | |
| Ever seriously considered committing suicide \$\frac{1}{2} \frac{1}{2} \frac{1} | |
| Seriously considered committing suicide in the past 12 months 2 th 12 months 2 th 12 months 2 th 12 months 3 th 14 th 12 months 3 th 14 th 13 months 3 th 14 th | |
| Seriously considered committing suicide but not in the past months \$\frac{2}{2} \frac{2}{2} \frac{2}{2 | |
| Not specified, seriously considered committing suicide in the 12 months \$^{2.12}\$ \$^{1.12}\$ \$^{1.12}\$ Never seriously considered committing suicide \$^{2.12}\$ Not specified, seriously considered committing suicide \$^{3.12}\$ \$^{1.14}\$ Not specified, seriously considered committing suicide \$^{3.12}\$ \$^{1.14}\$ Second perceived mental health \$^{4.2}\$ \$^{1.12}\$ Second perceived mental health \$^{4.2}\$ \$^{1.12}\$ Perc | IS |
| Not specified, seriously considered committing suicide \$\frac{1.11}{1.12}\$ \text{ Mot specified, seriously considered committing suicide \$\frac{1.11}{1.12}\$ \text{ Mot specified, seriously considered committing suicide \$\frac{1.11}{1.12}\$ \text{ Excellent or very good perceived mental health \$\frac{1.2}{1.12}\$ \text{ Good perceived mental health \$\frac{1.2}{1.12}\$ \text{ Perceived mental health \$\frac{1.2}{1.12}\$ \text{ Perceived mental health \$\frac{1.2}{1.12}\$ \text{ Perceived mental health not specified \$\frac{1.2}{1.12}\$ \text{ Perceived mental health not specified \$\frac{1.2}{1.12}\$ \text{ Seriously considered committing suicide in the past 12 months \$\frac{1.11}{1.12}\$ \text{ Seriously considered committing suicide but not in the past months \$\frac{1.11}{1.12}\$ \text{ Not specified, seriously considered committing suicide in the 12 months \$\frac{1.12}{1.12}\$ \text{ Not specified, seriously considered committing suicide \$\frac{1.12}{1.12}\$ Not specified, | 2 |
| Not specified, seriously considered committing suicide \$\frac{a}{2} \cdot \frac{1a}{2} \c | past |
| Total, perceived mental health 4.2.10 Excellent or very good perceived mental health 4.2.10 Good perceived mental health 4.2.11 Fair or poor perceived mental health 4.2.12 Perceived mental health not specified 4.2.12 Total, suicidal thoughts 5.14 Ever seriously considered committing suicide in the past 12 monts 5.14.13 Seriously considered committing suicide but not in the past months 5.14.13 Not specified, seriously considered committing suicide in the 12 months 5.14.13 Never seriously considered committing suicide in the 12 months 5.14.13 Never seriously considered committing suicide 5.15.14 Not specified, seriously considered committing suicide 5.15.14 Not specified, seriously considered committing suicide 5.15.14 Not specified, seriously considered committing suicide 5.15.14 Excellent or very good perceived mental health 4.2.10 | |
| Excellent or very good perceived mental health \$\frac{4}{2}\$ \$\frac{11}{12}\$ Good perceived mental health \$\frac{4}{2}\$ \$\frac{11}{12}\$ Fair or poor perceived mental health \$\frac{4}{2}\$ \$\frac{11}{12}\$ Perceived mental health not specified \$\frac{4}{2}\$ \$\frac{11}{2}\$ Total, suicidal thoughts \$\frac{5}{2}\$ \$\frac{14}{2}\$ Ever seriously considered committing suicide \$\frac{5}{2}\$ \$\frac{14}{2}\$ \$\frac{15}{2}\$ Seriously considered committing suicide but not in the past months \$\frac{5}{2}\$ \$\frac{14}{2}\$ \$\frac{13}{2}\$ Not specified, seriously considered committing suicide in the 12 months \$\frac{5}{2}\$ \$\frac{14}{2}\$ \$\frac{15}{2}\$ \$\fr | |
| Good perceived mental health \$\frac{4}{2}\$.\$\frac{12}{12}\$ Perceived mental health not specified \$\frac{4}{2}\$.\$\frac{12}{12}\$ Perceived mental health not specified \$\frac{4}{2}\$.\$\frac{12}{12}\$ Total, suicidal thoughts \$\frac{5}{2}\$.\$\frac{14}{12}\$ Both sexes Ever seriously considered committing suicide \$\frac{5}{2}\$.\$\frac{14}{12}\$ Seriously considered committing suicide in the past 12 months \$\frac{5}{2}\$.\$\frac{14}{12}\$ Not specified, seriously considered committing suicide in the 12 months \$\frac{5}{2}\$.\$\frac{14}{12}\$ Never seriously considered committing suicide \$\frac{5}{2}\$.\$\frac{14}{2}\$ Never seriously considered committing suicide \$\frac{5}{2}\$.\$\frac{14}{2}\$ Not specified, seriously considered committing suicide \$\frac{5}{2}\$.\$\frac{14}{2}\$ Excellent or very good perceived mental health \$\frac{4}{2}\$.\$\frac{12}{2}\$ | |
| Fair or poor perceived mental health 4 2 12 Perceived mental health not specified 4 2 13 Total, suicidal thoughts 5 14 Ever seriously considered committing suicide 5 14 15 Seriously considered committing suicide in the past 12 months 5 14 15 Seriously considered committing suicide but not in the past months 5 14 15 Not specified, seriously considered committing suicide in the 12 months 5 14 15 15 Never seriously considered committing suicide 5 14 Never seriously considered committing suicide 5 12 14 Not specified, seriously considered committing suicide 5 12 14 Not specified, seriously considered committing suicide 5 12 14 Excellent or very good perceived mental health 4 2 18 | |
| Fair or poor perceived mental health 4.2.12 Perceived mental health not specified 4.2.13 Total, suicidal thoughts 2.14 Ever seriously considered committing suicide 2.14.15 Seriously considered committing suicide in the past 12 months 2.14.15 Seriously considered committing suicide but not in the past months 2.14.15 Not specified, seriously considered committing suicide in the 12 months 2.14.15 Never seriously considered committing suicide 5.12 Never seriously considered committing suicide 5.12 Not specified, seriously considered committing suicide 5.12.14 Not specified, seriously considered committing suicide 5.12.14 Excellent or very good perceived mental health 4.2.16 | |
| Perceived mental health not specified 4-2-11 Total, suicidal thoughts 5-14 Ever seriously considered committing suicide 5-14-15 Seriously considered committing suicide in the past 12 mon 5-14-15 Seriously considered committing suicide but not in the past months 5-14-15 Not specified, seriously considered committing suicide in the 12 months 5-14-15-15 Never seriously considered committing suicide 5-14 Not specified, seriously considered committing suicide 5-14 Not specified, seriously considered committing suicide 5-13-14 Not specified, seriously considered committing suicide 5-13-14 Excellent or very good perceived mental health 4-2-10 | |
| Both sexes Ever seriously considered committing suicide \$\frac{5}{14}\$ 15 Seriously considered committing suicide in the past 12 mon \$\frac{5}{14}\$ 15 Seriously considered committing suicide but not in the past months \$\frac{5}{14}\$ 15 Not specified, seriously considered committing suicide in the 12 months \$\frac{5}{14}\$ 15 Never seriously considered committing suicide \$\frac{5}{14}\$ 15 Not specified, seriously considered committing suicide \$\frac{5}{14}\$ 15 Not specified, seriously considered committing suicide \$\frac{5}{14}\$ 15 Not specified, seriously considered committing suicide \$\frac{5}{5}\$ 15 Male Total, perceived mental health \$\frac{4}{5}\$ 2 Excellent or very good perceived mental health \$\frac{4}{5}\$ 2.10 | |
| Ever seriously considered committing suicide 5, 14, 15 Seriously considered committing suicide in the past 12 mon so 14, 15 Seriously considered committing suicide but not in the past months 5, 14, 15 Not specified, seriously considered committing suicide in the 12 months 5, 13, 14, 15 Never seriously considered committing suicide 5, 14 Not specified, seriously considered committing suicide 5, 13, 14 Not specified, seriously considered committing suicide 5, 13, 14 Not specified, seriously considered committing suicide 5, 13, 14 Excellent or very good perceived mental health 4, 2, 10 | |
| Seriously considered committing suicide in the past 12 months 5 14 15 Not specified, seriously considered committing suicide but not in the past months 5 14 15 Never seriously considered committing suicide in the 12 months 5 14 14 12 Never seriously considered committing suicide 5 14 Not specified, seriously considered committing suicide 5 13 14 Not specified, seriously considered committing suicide 5 13 14 Excellent or very good perceived mental health 4 2 10 | |
| Seriously considered committing suicide but not in the past months \$\frac{5}{2}\$ \$\frac{14}{15}\$\$ Not specified, seriously considered committing suicide in the 12 months \$\frac{5}{2}\$ \$\frac{14}{15}\$ \$\frac{12}{15}\$\$ Never seriously considered committing suicide \$\frac{5}{2}\$ \$\frac{14}{15}\$\$ Not specified, seriously considered committing suicide \$\frac{5}{2}\$ \$\frac{13}{15}\$ \$\frac{14}{15}\$\$ Male Total, perceived mental health \$\frac{4}{2}\$ \$\frac{2}{15}\$ \$\frac{14}{15}\$\$ Excellent or very good perceived mental health \$\frac{4}{2}\$ \$\frac{10}{15}\$ | |
| Not specified, seriously considered committing suicide in the 12 months \$\frac{5}{2} \frac{14}{2} \frac{12}{2}\$ Never seriously considered committing suicide \$\frac{5}{2} \frac{14}{2}\$ Not specified, seriously considered committing suicide \$\frac{5}{2} \frac{14}{2}\$ Male Total, perceived mental health \$\frac{4}{2} \frac{2}{2}\$ Excellent or very good perceived mental health \$\frac{4}{2} \frac{10}{2}\$ | |
| Never seriously considered committing suicide ^{5, 14} Not specified, seriously considered committing suicide ^{5, 12, 14} Male Total, perceived mental health ^{4, 2} Excellent or very good perceived mental health ^{4, 2, 10} | 2 |
| Not specified, seriously considered committing suicide 5, 13, 14 Male Total, perceived mental health 4, 2 Excellent or very good perceived mental health 4, 2, 10 | past |
| Male Total, perceived mental health 4.2 Excellent or very good perceived mental health 4.2.10 | |
| Excellent or very good perceived mental health 4: 2: 10 | |
| | |
| Good perceived mental health 4-2-11 | |
| | |
| Fair or poor perceived mental health 4: 2: 12 | |
| Perceived mental health not specified ^{4,2,13} | |
| Total, suicidal thoughts ^{5, 14} | |
| Ever seriously considered committing suicide 5, 14, 15 | |

| Aboriginal identity ⁶ | Age group | Sex | Perceived mental health and suicidal thoughts | 2012 |
|----------------------------------|-----------|--------|---|------|
| | | | Seriously considered committing suicide in the past 12 months 5.14.15 | |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 15, 15} | 1 |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 14, 15} | |
| | | | Never seriously considered committing suicide 5. 14 | |
| | | | Not specified, seriously considered committing suicide ^{5, 13, 14} | |
| | | | Total, perceived mental health 4· 2 | |
| | | | Excellent or very good perceived mental health 4.2.10 | |
| | | | Good perceived mental health 4.2.11 | |
| | | | Fair or poor perceived mental health 4. 2. 12 | |
| | | | Perceived mental health not specified 4, 2, 13 | |
| | | | Total, suicidal thoughts 5.14 | |
| | | Female | Ever seriously considered committing suicide 5, 14, 15 | |
| | | | Seriously considered committing suicide in the past 12 months 5 14 15 | |
| | | | Seriously considered committing suicide but not in the past 12 months ^{5, 14, 15} | |
| | | | Not specified, seriously considered committing suicide in the past 12 months ^{5, 13, 14, 15} | |
| | | | Never seriously considered committing suicide ^{5, 14} | |
| | | | Not specified, seriously considered committing suicide 5, 13, 14 | |

Symbol legend:

Use with caution

Too unreliable to be published

Footnotes:

- Source: Aboriginal Peoples Survey 2012.
- This table was previously released March 30, 2015. This current version of the table has been modified from the original version by (1) the addition of provincial and territorial data, (2) the recalculation of some percentages, based on the overall total for the variable (see footnotes 9 and 14), and (3) the change of the theme "Perceived mental health and suicidal thoughts and attempts" to "Perceived mental health and suicidal thoughts". Sub-categories of "Seriously considered committing suicide in the past 12 months", which indicated whether or not the person had ever attempted suicide, have been removed.

Back to original table

Date modified: 2015-07-14

- Excludes persons living on Indian reserves or settlements.

 Perceived mental health refers to the perception of a person's mental health in general. The question on perceived mental health was only asked of respondents who completed their own (non-proxy) interviews. Respondents were asked "In general, would you say your mental health is... Excellent? Very good? Good? Fair?
- Questions on suicidal thoughts were only asked of respondents who completed their own (non-proxy) interviews. Respondents were asked "Have you ever seriously
- considered committing suicide or taking your own life?".

 Aboriginal identity includes persons living off reserve who reported being an Aboriginal person, that is, First Nations (North American Indian), Métis or Inuk (Inuit) and/or those who reported Registered or Treaty Indian status, that is registered under the Indian Act of Canada, and/or those who reported membership in a First Nation or Indian band. The sum of the categories included in this variable is greater than the total population estimate for Aboriginal identity because a person may have reported more than one Aboriginal identity: for example, a person could self-identify as both First Nations and Métis.
- Atlantic includes Newfoundland and Labrador, Prince Edward Island, Nova Scotia, and New Brunswick.
- Registered or Treaty Indian status refers to whether or not a person reported being a Registered or Treaty Indian. Registered Indians are persons who are registered under the Indian Act of Canada. Treaty Indians are persons who belong to a First Nation or Indian band that signed a treaty with the Crown. Registered or Treaty Indians are sometimes also called Status Indians.
- All percentages for the component "perceived mental health" are calculated using the number of persons in "total, perceived mental health" as denominator. This category includes persons who rated their mental health as "excellent" or "very good". These are the most positive ratings on the perceived mental health
- scale, a five-point scale ranging from "excellent" to "poor"
- This category includes persons who rated their mental health as "good". This is the midpoint on the perceived mental health scale which is a five-point scale ranging 11. from "excellent" to "poor"
- 12. This category includes persons who rated their mental health as "fair" or "poor". These are the most negative ratings on the perceived mental health scale, a fivepoint scale ranging from "excellent" to "poor".
 "Not specified" categories consist of item non-response ("Refusal", "Don't know", and "Not stated").
- All percentages for the component "suicidal thoughts" are calculated using the number of persons in "total, suicidal thoughts" as denominator.
- Respondents who said they had ever seriously considered committing suicide or taking their own life were asked: "Has this happened in the past 12 months?". Bootstrapping techniques were used to produce the coefficient of variation (CV). 15.
- Data with a coefficient of variation (CV) from 16.6% to 33.3% are identified by an (E) and should be interpreted with caution. 17.
- Data with a coefficient of variation (CV) greater than 33.3% were suppressed (F) due to extreme sampling variability

Source: Statistics Canada. Table 577-0013 - Aboriginal peoples survey, perceived mental health and suicidal thoughts, by Aboriginal identity, age group and sex, population aged 18 years and over, Canada, provinces and territories, occasional, CANSIM (database). (accessed:)