

# Suicide (NHC) - deaths per 100 000 persons, age standardised

Geography	Nordland						Troms						Finnmark					
	men			women			men			women			men			women		
Age	0-44 yrs	45-74 yrs	75 yrs +	0-44 yrs	45-74 yrs	75 yrs +	0-44 yrs	45-74 yrs	75 yrs +	0-44 yrs	45-74 yrs	75 yrs +	0-44 yrs	45-74 yrs	75 yrs +	0-44 yrs	45-74 yrs	75 yrs +
Year																		
1970-1974	7	15	-	3	3	-	8	18	-	4	10	-	7	20	-	-	-	-
1971-1975	6	17	-	3	2	-	8	17	-	4	11	-	5	26	-	3	-	-
1972-1976	6	17	-	3	2	-	8	20	-	3	13	-	5	26	-	3	-	-
1973-1977	9	15	13	3	3	-	9	23	-	4	13	-	6	31	-	3	-	-
1974-1978	9	14	11	3	2	-	9	19	-	5	12	-	6	30	-	3	-	-
1975-1979	8	14	11	4	3	-	8	25	-	5	15	-	7	26	-	4	-	-
1976-1980	8	21	14	4	4	-	9	27	-	4	11	-	7	31	-	2	-	-
1977-1981	11	21	13	4	5	-	9	26	-	5	9	-	10	42	-	2	-	-
1978-1982	10	29	9	5	7	-	13	23	-	4	10	-	9	36	-	6	-	-
1979-1983	12	31	12	5	9	-	15	26	-	3	9	-	15	34	-	6	-	-
1980-1984	15	36	18	4	9	-	16	24	-	3	6	-	24	28	-	5	-	-
1981-1985	16	32	14	5	12	-	17	19	-	3	4	-	33	20	-	6	-	-
1982-1986	15	32	14	4	12	-	20	18	-	3	8	-	35	10	-	6	-	-
1983-1987	16	26	31	4	11	-	20	23	-	4	10	-	41	12	-	2	-	-
1984-1988	16	26	27	4	9	-	22	18	-	3	10	-	42	20	-	4	-	-
1985-1989	16	22	31	5	9	-	22	25	-	3	11	-	36	28	-	4	-	-
1986-1990	17	27	33	4	10	-	21	30	-	4	13	-	30	28	-	4	-	-
1987-1991	17	30	33	5	11	-	19	32	-	4	12	-	30	29	-	6	-	-
1988-1992	17	31	18	6	12	-	16	30	-	2	9	-	27	33	-	9	-	-
1989-1993	18	31	32	5	12	-	17	32	-	4	8	-	27	25	-	7	-	-
1990-1994	18	33	33	5	12	:	17	27	:	5	7	:	31	19	:	8	:	:
1991-1995	18	26	33	5	9	:	17	24	:	4	7	:	30	23	:	9	:	:
1992-1996	19	24	35	3	9	:	18	24	:	4	8	:	30	28	:	9	:	:
1993-1997	19	23	34	4	8	:	17	26	:	4	9	:	28	28	:	:	:	:
1994-1998	18	22	22	4	10	:	15	26	:	4	9	:	26	34	:	:	:	:
1995-1999	17	19	22	3	10	:	14	26	:	5	10	:	21	44	:	:	:	:
1996-2000	15	21	20	3	9	:	15	24	:	6	8	:	24	40	:	:	:	:
1997-2001	15	22	26	4	8	:	13	23	:	6	4	:	25	41	:	9	:	:
1998-2002	15	22	30	4	8	:	13	19	:	6	3	:	25	37	:	7	:	:
1999-2003	15	21	27	6	8	:	12	19	:	6	3	:	23	32	:	:	:	:
2000-2004	16	22	26	6	7	:	13	17	:	5	3	:	28	23	:	:	:	:
2001-2005	16	19	26	6	7	:	13	16	:	6	7	:	29	20	:	7	:	:
2002-2006	14	18	21	6	8	:	15	13	:	6	8	:	28	17	:	5	:	:
2003-2007	13	15	15	5	7	:	15	15	:	5	9	:	25	17	:	4	:	:
2004-2008	14	16	15	5	6	:	14	16	:	6	9	:	26	15	:	6	:	:
2005-2009	14	16	8	4	5	:	14	20	:	7	8	:	25	16	:	8	:	:
2006-2010	13	18	10	4	5	:	13	18	:	5	6	:	23	24	:	8	:	:
2007-2011	16	17	12	4	5	:	12	19	:	6	6	:	19	24	:	9	:	:

# Suicide (NHC) - *deaths per 100 000 persons, age standardised*

Geography	Nordland						Troms						Finnmark					
	men			women			men			women			men			women		
Age	0-44 yrs	45-74 yrs	75 yrs +	0-44 yrs	45-74 yrs	75 yrs +	0-44 yrs	45-74 yrs	75 yrs +	0-44 yrs	45-74 yrs	75 yrs +	0-44 yrs	45-74 yrs	75 yrs +	0-44 yrs	45-74 yrs	75 yrs +
Year																		
2008-2012	19	18	20	4	5	:	12	15	:	6	7	:	23	21	:	10	:	:
2009-2013	19	18	19	5	6	:	12	16	:	5	8	:	20	19	:	7	:	:
2010-2014	18	17	29	5	6	:	12	11	:	3	7	:	15	19	:	5	:	:
2011-2015	19	17	29	5	6	:	12	11	:	2	7	:	11	16	:	7	:	:
2012-2016	17	17	32	6	5	:	13	12	:	2	7	:	10	17	:	:	:	:
2013-2017	16	19	35	7	6	:	12	14	:	3	7	:	10	15	:	:	:	:

## CellMark Legend

- .. Missing data
  - .
  - Not possible to calculate
  - :
  - Hidden value
- obs: Number of deaths is only shown at country level.

## Description

Suicide and intentional self-harm, including sequelae of intentional self-harm (ICD-10: X60-X84, Y87.0). Number of deaths per 100 000 inhabitants\* per year, standardized for age composition. Five-year moving averages, i.e. averages for overlapping five-year periods.

To change the table, open "change selection of...".

Four measures are available. Use the Measure button to select:

1. Number of deaths (Only shown for Norway, for privacy reasons)
2. Deaths per 100 000 persons = Number per 100 000 inhabitants\* per year (rate)
3. Per 100 000, standardized = Number per 100 000 inhabitants\* per year (rate). Standardized for age composition.
4. Ratio (Norway=100) = Ratio between the county's standardized rate and the national rate for a given year, presented as five-year average. Examples; ratio = 130 means that the county's standardized rate is 30% higher than the national level. A ratio of 87 means that the county's rate is 13% lower than the national level.

\*Includes residents in the relevant age and gender segment.

The statistics include all persons registered as a resident of Norway at the time of death, regardless of whether the death occurred in Norway or abroad.

Rates are age standardised to reduce the effect of differing age distributions when groups are compared over time and between geographical regions.

Annual figures are standardised according to 5-year age groups in the Norwegian population per 1 January 2012.

## Source

Cause of Death Registry, Norwegian Institute of Public Health

## Collection

Based on information from the Cause of Death Registry. The Norwegian Institute of Public Health is responsible for the registry.

Mortality statistics are collated from death reports completed by public physicians. Additional information is gathered from results of autopsies and forensic examinations. Cause of death is coded according to the underlying cause of death cited by the physician who signs the death certificate. Underlying causes of death are defined as the illness or injury that, in the opinion of the physician, started the chain of conditions leading directly to death. Causes of death are coded according to the principles and directions of the World Health Organisation (WHO). The international classification system ICD is used. The following editions of the ICD system have been used: 1970-1985: ICD-8, 1986-1995: ICD-9, 1996 onwards: ICD-10.

## Interpretation and sources of error

Sequelae of intentional self-harm (ICD-10: Y87.0) is included with the definition of suicide and intentional self-harm.

## Data quality

Good.

## When numbers are missing

Statistics based on fewer than 3 cases is hidden to protect the identity of the persons. Figures are also hidden when the population group in which the cases originate, is smaller than 10 persons. If more than 20 per cent of the cases in a time series are hidden due to the abovementioned reasons, the complete time series is hidden to avoid giving a skewed impression.

Some values are hidden to prevent their use in calculating a value hidden for privacy reasons.

The entire time series is removed if over half the rates are based on 6 or fewer deaths. This is because of reduced reliability. This rule is applied separately to the time periods 1970-1989 and from 1990.

# Suicide (NHC) - *deaths per 100 000 persons, age standardised*

## Time periods

1970-1974 to 2013-2017 (5-year moving averages)

## Geographical level

Country, health region, counties

## Gender

Men, women.

## Age groups

Age at death: All ages, 0-74 years, 0-44 years, 0-14 years, 15-24 years, 25-44 years, 45-74 years, 75 years and older

## Frequency of updates

Annually

## Last updated

10/12/18

## Keywords

Click on a keyword to search for similar indicators.

- Cause of death
- Dead
- Death
- Diagnosis
- External cause
- Suicide
- Violent death

## Fact sheets

Below are links to relevant fact sheets, articles and reports. These may describe trends over time in the Norwegian population or differences by sex, age group, geographical region or socioeconomic status:

- [Suicide and suicide attempts in Norway - fact sheet](#)
- [Suicide prevention - fact sheet \(in Norwegian only\)](#)